

- b*¹³. Protarsi i. and ii. with a central apical spine beneath.
- a*¹⁴. Eyes in two rows AGELENIDÆ (p. 332).
- b*¹⁴. Eyes in three rows PISAURIDÆ (p. 303).
- b*¹². Tibia of male palpus without an apical external spur.
- a*¹⁵. Eyes of anterior row subequal or centrals sometimes larger. Labium and maxillæ shorter in proportion LYCOSIDÆ (p. 314).
- b*¹⁵. Central eyes of anterior row much smaller. Labium and maxillæ longer in proportion OXYOPIDÆ (p. 338).
- II. Tarsus iv. set with numerous serrate bristles, or with a single series of serrate spines beneath.
- a*¹⁶. Tarsus iv. set with a single series of stout, curved, serrate spines beneath THERIDIIDÆ (p. 371).
- b*¹⁶. Tarsus iv. set with numerous serrate bristles.
- a*¹⁷. Tibiæ and protarsi i. and ii. furnished with a series of widely separate, very long spines, the interspaces being set with a row of shorter curved spines, increasing in length distally MIMETIDÆ (p. 409).
- b*¹⁷. Tibiæ and protarsi not so furnished.
- a*¹⁸. Legs shorter and stouter. Maxillæ not inclined over labium ARGYOPIDÆ (p. 412).
- b*¹⁸. Legs very long and slender. Maxillæ inclined over labium PHOLCIDÆ (p. 365).
- bb*. Eyes 6.
- a*¹⁹. Mandibles not soldered together at their base. Fang of mandible long and slender LEPTONETIDÆ (p. 45).
- b*¹⁹. Mandibles soldered together at their base. Fang of mandible short, but weak. Colulus present or absent SCYTODIDÆ (p. 48).

The following genera or species have been added to the Mexican or Central-American fauna during the course of publication of the present volume, or have been accidentally omitted in our enumeration:—

THERAPHOSIDÆ.

- Citharacanthus*, Pocock, Ann. & Mag. Nat. Hist. (7) viii. p. 551 (1901), n. g., for *Eurypelma longipes*, F. Cambr. antea, p. 21.
- Dugesiella crinita*, Pocock, loc. cit. p. 552.—Mexico, Guanajuato.
- Aphonopelma*, Pocock, loc. cit. p. 553, n. g., for *Eurypelma seemanni*, F. Cambr. antea, p. 26.