

Colour. Body, including the underside, coxae of legs, mandibles, and palpi, dull orange; eye-tumulus black; upperside mottled with brown; the second and third dorsal opisthosomatic segments with a subcentral yellow mark on each side, longitudinally; legs jet-black, protarsi and tarsi piceous-brown.

Hab. MEXICO, Omilteme in Guerrero (*H. H. Smith*).

The following species of Phalangiids, probably belonging to the group "Plagiostethi," are unknown to me:—

Egænus mexicanus, Becker, C. R. Ent. Belg. (3) no. 69, p. xxvi.

Hab. MEXICO.

Opilio ischionotatus, Dugès, La Nat. Mex. vii. p. 194, t. 2. figg. 1-10.

Hab. MEXICO, Guanajuato.

Liobunum mexicanum, Banks, Journ. N. Y. Ent. Soc. vi. p. 181 (1898).

Hab. MEXICO, Amecameca (*Barrett*).

Liobunum albipalpe, Banks, Journ. N. Y. Ent. Soc. vi. p. 181 (1898).

Hab. MEXICO.

Liobunum denticulatum, Banks, Journ. N. Y. Ent. Soc. viii. p. 199 (1900).

Hab. MEXICO, Cuernavaca, Morelos (*Barrett*).

Liobunum consimile, Banks, Journ. N. Y. Ent. Soc. viii. p. 200.

Hab. MEXICO, Cuernavaca, Morelos (*Barrett*).

Leptobunus spinulatus, Banks, Journ. N. Y. Ent. Soc. vi. p. 182.

Hab. MEXICO.

Hoplobunus barretti, Banks, Journ. N. Y. Ent. Soc. viii. p. 200.

Hab. MEXICO, Cuernavaca, Morelos (*Barrett*).

A D D E N D A.

Prionostemma scintillans, Pocock, Ann. & Mag. Nat. Hist. (7) xi. p. 434 (1903).

Hab. GUATEMALA, Barrancos (*Stoll*).

Prionostemma bicolor, Pocock, loc. cit. p. 434.

Hab. GUATEMALA, Barrancos (*Stoll*).