

- Elytra yellow, each with two spots at the base, a median fascia, and the apex black *mylabrina*, Chevr.
- Elytra yellow, each with a large spot at the base and a very broad common post-median fascia black *rugulipennis*, n. sp. ✓
- Elytra yellow, each with two spots at the base and a broad more or less elongate post-median fascia, the latter not reaching the suture, and sometimes confluent with the humeral spot, black *postica*, Lec. ✓
- Elytra yellow or reddish-yellow, each with a median vitta, sometimes narrowed in front and with a spot separated from it at the base on the inner side, black.
- Suture pale.
- Surface vaguely sulcate *tenuicostatis*, Dugès. ✓
- Surface not sulcate *divirgata*, Vill. & Peñaf. ✓
- Suture piceous or black, usually more broadly so towards the base *decorata*, Haag. ✓
- Elytra black, with the sutural, apical, and lateral margins reddish-yellow *akhurstiana*, Horn. ✓
- Elytra black, with the sutural and lateral margins narrowly, an oblique basal stripe on the disc, and a subapical streak, yellow *insulata*, Lec. ✓
- Elytra yellow, each with two narrow black vittæ *bilineata*, Horn. ✓
- Elytra unicolorous, luteous or reddish-brown.
- Surface sulcate.
- Thorax thickly punctured *hirticollis*, n. sp. ✓
- Thorax very sparsely punctured *mariarum*, n. sp.
- Surface not sulcate.
- Antennæ rather slender; thorax with a black cruciform mark on the disc *nobilis*, Haag.
- Antennæ shorter, stout, the joints ovate; thorax with two black spots on the disc *quadrinervata*, Herr. & Mend.

1. *Pyrota terminata*. (Tab. XX. fig. 1, ♂.)

Pyrota terminata, Lec. New Sp. Col. p. 159 (1866)¹; Horn, Trans. Am. Ent. Soc. xii. p. 113 (nec Sturm)².

Lytta mylabrina, Lec. Proc. Ac. Phil. vi. p. 337³.

Pyrota mylabrina, Horn, Trans. Am. Ent. Soc. xii. p. 112, t. 5. fig. 13 (nec Chevr.)⁴.

Pyrota concinna, Casey, Ann. N. York Acad. vi. p. 174 (1891)⁵.

Hab. NORTH AMERICA, Kansas¹, Colorado⁴, Missouri², Western Texas⁵, Mexican boundary³.—MEXICO⁴, Villa Lerdo in Durango, Chihuahua city, Monterey in Nuevo Leon (*Höge*).

Found in plenty by Herr Höge at Villa Lerdo. With the numerous Mexican and Texan examples before me I am unable to distinguish more than one species.

The type of *P. terminata* has entirely black legs; in all the specimens I have seen