

*Hab.* PANAMA, Chiriqui (*Zahn*), Bugaba (*Arcé*<sup>1</sup>).

This form, which is very closely allied to *A. miltiades*, differing from it chiefly in the absence of the discal white spot of the secondaries, appears to be restricted to Chiriqui and its vicinity, where it exists to the exclusion of the allied form.

### 3. *Antirrhæa pterocopa*. (Tab. VII. figg. 4, 5, 6.)

*Antirrhæa pterocopa*, Salv. & Godm. Ann. & Mag. N. H. ser. 4, ii. p. 143.

♂ alis anticis fuscis, punctis tribus minutis griseis in area ultra cellulam positis; posticis obscurioribus, macula fulva irregulari ad apicem, puncto minuto griseo infra eam, maculis tribus cyaneis serie alam transeunte: subtus alis griseis fusco irroratis, fascia communi obscura arcuata a costa anticarum ad angulum analem posticarum extendente alteraque subparallela interiore griseo-purpureo extus marginatis, serie irregulari punctorum submarginalium albescentium punctoque nigro basin posticarum versus notatis.

♀ mari similis, sed colore fusco dilutiore, anticarum fascia transversa ultra cellulam cyanea notata maculas albidas includente, plaga fulva posticarum multo majore et margine externo late fulvo notato, maculis posticarum cyaneis ad marginem internum absentibus.

*Hab.* COSTA RICA (*Endres*); PANAMA, Santa Fé and Calobre (*Arcé*).

The specimen from which our original description was taken is a male in bad condition, sent by our collector, Arcé, from Santa Fé; since then we have received others of both sexes from the same country and also from Costa Rica. In colour it most resembles *A. philopæmen*, from the Upper-Amazon region; but it differs very considerably in the form of its anterior wings. These at their apex are abruptly obtuse instead of rounded, and the outer margin strongly angulated at the extremity of the upper radial. The great difference, too, in the size and coloration of the sexes is remarkable, no such contrast being seen in *A. philopæmen*. It appears to be a scarce insect, as we have received but few examples of it, notwithstanding the large collections that have come to us from the country it inhabits.

## TISIPHONE.

*Tisiphone*, Hübner, Samml. ex. Schmett. ii. t. 86. (Type *T. hercyna*, Hübn.)

*Tisiphone*, Westwood, Gen. Diurn. Lep. ii. p. 370; Butl. Cat. Sat. B. M. p. 71.

When first used by Hübner\* this genus included three species, none of which belongs to *Tisiphone* as now understood. He afterwards added *T. hercyna*; and this has long stood as the only representative of the genus, to the exclusion of the other three. According to strict law the modern practice is incorrect; but in dealing with Hübnerian genera great inconvenience and confusion would arise by enforcing conformity, and no doubt *Tisiphone* had best be left to stand as the name of this genus.

One of the distinguishing characters between *Tisiphone* and *Euptychia* and *Taygetis*

\* Verz. bek. Schm. p. 60.