

Hab. MEXICO, Huahuapam (*Hedemann*¹), Cuernavaca (*Mus. Felder*.¹); GUATEMALA, Choctum (*Hague*³).

The eyes in this species are destitute of hairs, and the palpi more densely hairy than in almost any other *Euptychia* that we have examined. These and other less important characters induced R. Felder to place *E. squamistriga* in a genus by itself; and we should be glad to adopt this division, were we able to trace any well-defined group of insects possessing such characteristics. But the hairiness or smoothness of the eyes, so far as we have been able to see, is subject to considerable variation in *Euptychia*, and still more so the hairiness of the palpi; so that we have not been able to make any satisfactory divisions resting upon these features. The neuration of the primaries places *E. squamistriga* with the insects which here precede it, where it may rest pending a thorough examination of the whole genus.

This species is apparently a rare one, as we have received but a single specimen from Guatemala, the type of *E. zabdi*, now figured. In Mexico, besides the type obtained by Lieut. v. Hedemann at Huahuapam, near Cordova, Felder speaks of two other males in his collection captured at Cuernavaca.

The names given by R. Felder and Mr. Butler to this butterfly were proposed within a few weeks of each other. The paper containing the description of the former author was read, but probably not published, prior to the issue of the part of 'Lepidoptera Exotica' containing the description and figure of the latter. As Mr. Butler in his corrections places his name, *E. zabdi*, as a synonym of *E. squamistriga*, we follow him in so doing.

b'. First subcostal branch of primaries emitted before the end of the cell, the second beyond it.

13. *Euptychia similis*. (Tab. VIII. fig. 4.)

Euptychia similis, Butl. P. Z. S. 1866, p. 463¹.

? *Euptychia similis*, var., Butl. P. Z. S. 1866, p. 463², 1867, t. 12. f. 10³.

? *Euptychia undina*, Butl. Ent. Month. Mag. vi. p. 252⁴.

E. labæ affinis, sed alis anticis subtus haud fulvo notatis et serie ocellorum submarginali obsoleta, ocello ad angulum analem tantum bene definito; striis transversis multo angustioribus.

Hab. MEXICO, Oaxaca (*Rébouch*); GUATEMALA, San Gerónimo (*Champion*), central valleys (*F. D. G. & O. S.*¹ & *H. Hague*); ? NICARAGUA (*Mus. Brit.*^{2 3 4}).

Though closely resembling *E. pieria*, this butterfly appears to be sufficiently distinct from that species to be recognizable. Besides the slight difference in the position of the second subcostal branch of the primaries, the chief differences observable are the position of the ocellus of the anal angle of the hind wings beneath (which in *E. similis*