

ad marginem internum (extus pallide griseo limbata) et ocellis quinque serie submarginali positis ornatis, primo maximo, reliquis obsoletis sed argenteo indicatis.

Hab. NORTH AMERICA, Texas¹.—MEXICO (*Deppe, Mus. Berol.*); GUATEMALA, Cachil, Vera Paz (*Champion*).

Mr. Champion has recently sent us several specimens of this pretty species, all taken 3000 or 4000 feet above the sea, on the steep hill-side near the village of Cachil, and overlooking the plain of Salamá in Vera Paz. These agree very fairly with Mr. W. H. Edwards's description of his *E. rubricata*, described from Texan specimens. In Guatemala the insect is doubtless very local in its distribution. Mr. Champion only obtained it in one place; and we never met with it ourselves, nor have we received it from any other collector. Two years ago we examined five specimens of a *Euptychia* in the Berlin Museum, two of which were of Deppe's collecting, the others from Texas. From our notes we have little doubt they all belong to *E. rubricata*.

c'. First and second subcostal branches of primaries emitted before the end of the cell.

16. *Euptychia oreba*.

Euptychia oreba, Butl. Cist. Ent. i. p. 19¹; Ent. Monthly Mag. vi. p. 252, t. 1. f. 7²; Dist. Pr. Ent. Soc. 1876, p. xiii³.

Alis supra ferrugineo-fuscis; posticis rotundatis, extus paulo sinuatis: subtus alis cinerascentioribus; linea submarginali nigricante, altera valde sinuata, tertia fere recta ultra cellulam, et quarta indistincta per cellulam notatis; anticis ocellis quinque albo pupillatis sed indistinctis serie submarginali positis; posticis quoque ocellis quinque notatis, secundo et tertio obsoletis.

Hab. COSTA RICA (*Gabb*³).—COLOMBIA.

This species was originally described from specimens of which the locality was not recorded¹. These are now before us, and agree fairly with the specimen referred by Mr. Distant to *E. oreba* in his supplementary enumeration of Costa-Rican butterflies³. We have also specimens from the neighbourhood of Santa Marta, Colombia, which we consider to belong to the same species. Between these specimens, five in number, a certain amount of variation exists, but not enough to justify their separation without more materials. The Costa-Rica example comes nearest to the type.

17. *Euptychia libye*.

Papilio libye, Linn. Syst. Nat. ii. p. 772¹.

Euptychia libye, Butl. P. Z. S. 1866, p. 486²; Butl. & Druce, P. Z. S. 1874, p. 336³.

Euptychia libyoidea, Butl. P. Z. S. 1866, p. 487⁴, 1867, pl. 11. f. 13⁵; Dist. Pr. Ent. Soc. 1876, p. xiii⁶.

Alis supra fuscis, marginibus externis lineis duabus angustis circumcinctis: subtus alis multo pallidioribus; posticis purpureo tinctis, fasciis duabus ferrugineis, una per cellulam, altera ultra eam transfasciatis, lineis duabus submarginalibus et plaga elongata intra eas notatis; anticis ocello parvo in apice notatis; posticis ocellis quinque ornatis, primo et quarto multo majoribus et bipupillatis, quinto parvo, reliquis obsoletis.