

Hab. GUATEMALA, Polochic valley (*F. D. G. & O. S.*²), Chisoy valley (*Hague*), Choctum (*Hague*), Retalhuleu (*F. D. G. & O. S.*), Cubilguitz, Chiacam, Cahabon and Zapote (*Champion*); NICARAGUA⁴, Chontales (*Belt, Janson*); COSTA RICA (*v. Patten*³, *Gabb*⁶); PANAMA, Chiriqui, Calobre, and Santa Fé (*Arcé*).—SOUTH AMERICA to North Brazil; JAMAICA.

This is one of the commonest species of *Euptychia* in Central America; and we have received numbers of specimens from Guatemala. It varies to some extent in the markings of the underside; but we fail, at present, to see any definite geographical races, though Mr. Butler has separated an insect of this form from Nicaragua as a distinct species. In Guatemala, as doubtless elsewhere, *E. libye* is an inhabitant of the low-lying hot country.

18. *Euptychia tiessa*.

Euptychia tiessa, Hew. Equat. Lep. p. 37¹; Butl. Lep. Exot. p. 48, t. 18. f. 4².

Alis supra rufo-fuscis, linea obscura submarginali notatis; posticis dentatis et ocello nigro rubro circumdato ornatis: subtus ut supra, alis lineis duabus transversis mediis ferrugineis, linea duplici submarginali ejusdem coloris, inter has fascia lata grisea centro ferrugineo notatis; anticis ad apicem uno, posticis quinque ocellis submarginalibus ornatis, primo (maximo), quarto et quinto pupillatis, reliquis obsoletis.

Hab. NICARAGUA, Chontales (*Belt, Janson*); COSTA RICA (*v. Patten*), Irazu (*Rogers*); PANAMA, Calobre (*Arcé*).—ECUADOR.

This species may readily be distinguished from all its allies by the rich ferruginous markings of the underside, and by the broad submarginal grey band with a central ferruginous streak, in which the ocelli are placed. We have several examples taken by Mr. Belt at Chontales, in Nicaragua—southward of which locality it would appear to be rather a scarce insect, as we have only two specimens from Costa Rica and one from Panama. These, however, all agree with one we have from Guadalquiza, in Ecuador, where the types were obtained¹.

19. *Euptychia satyrina*. (Tab. VIII. fig. 23.)

Tagetis satyrina, Bates, Ent. Monthl. Mag. i. p. 179¹; Butl. & Druce, P. Z. S. 1874, p. 335².

Euptychia gigas, Butl. P. Z. S. 1866, p. 486, t. 40. f. 7³; Dist. Pr. Ent. Soc. 1876, p. xiii⁴.

Taygetis incerta, Butl. & Druce, Cist. Ent. i. p. 98⁵; Butl. Lep. Exot. p. 149, t. 53. f. 1⁶.

Alis fuscis (aliquando ferrugineo notatis); posticis ocello nigro albo pupillato, plerumque ferrugineo circumeincto (aliquando fere obsoleto), angulum analem versus: subtus dilutior, alis linea submarginali valde undulata, ferruginea, altera curvata interiore, tertia basin propiore; anticis punctis nigris indistinctis apud apicem; posticis ocellis quinque notatis, primo, quarto et quinto maximis.

Hab. MEXICO (*Mus. Brit.*); GUATEMALA, Polochic valley (*F. G. & O. S.*¹), Chuacus, Plain of Salamá, Choctum, Chisoy valley (*Hague*); Coban, Pancina, and San Gerónimo (*Champion*); COSTA RICA (*v. Patten*^{2 3 5}, *Gabb*⁴), Irazu (*Rogers*); PANAMA, Chiriqui (*Arcé*).