

24. **Euptychia phares.** (Tab. VIII. figg. 8, 9.)

Satyrus phares, Godt. Enc. Méth. ix. p. 491¹.

Neonympha phares, Westw. & Hew. Gen. Diurn. Lep. p. 375, t. 67. f. 4².

Euptychia phares, Butl. P. Z. S. 1866, p. 477³.

Alis supra fuscis, lineolis tribus subparallelis marginatis; anticis ocello unico in angulo apicali, posticis ocellis quatuor linea submarginali positis ornatis, primo et quarto maximis: subtus multo pallidioribus, lineis et ocellis multo magis distinctis, linea communi fusca extus sordide albido limbata et anticis plaga magna media fulva notatis.

Hab. PANAMA, Calobre (*Arcé*).—VENEZUELA³; BRAZIL¹.

Of this species we have a single female specimen, sent us from Calobre by Arcé, that now described and figured. This agrees closely with a series of examples from Brazil and Venezuela, showing that this insect, though apparently nowhere common, has a very extensive range.

25. **Euptychia harmonia.**

Euptychia harmonia, Butl. P. Z. S. 1866, p. 478, t. 39. f. 17¹.

Alis supra fuscis: subtus dilutioribus, lineolis tribus fuscis marginatis, fascia rufescens-fusca communi in posticis trisinuata extra cellulas, altera per cellulas transeunte; anticis ocellis quatuor vel quinque ornatis, eo ad angulum apicalem majore, reliquis obsoletis; posticis ocellis quinque notatis, quarto maximo, secundo quoad magnitudinem proximo, reliquis fere obsoletis, sed omnibus albo pupillatis.

Hab. COSTA RICA, Irazu and Caché (*Rogers*); PANAMA, Chiriquí (*Ribbe*).—ECUADOR¹.

This species, which by its neuration belongs to the same group as *E. camerta*, was originally described from Ecuadorean specimens¹. Since then we have received a number of examples both from Costa Rica and from the neighbourhood of Chiriquí. These all agree accurately with the typical form from Ecuador.

f. Wings, especially the apical portion of the secondaries, more or less washed with shining blue in both sexes; first and second subcostal branches emitted before the end of the cell; middle discocellular angular, and emitting a short recurrent nervule.

26. **Euptychia ebusa.**

Papilio ebusa, Cram. Pap. Ex. iv. t. 292. f. F, G¹.

Euptychia ebusa, Butl. P. Z. S. 1866, p. 489².

Euptychia arnæa, Butl. & Druce, P. Z. S. 1874, p. 337 (ex Fabr.)³.

♂ alis supra pallide fuscis, lineis paginæ inferioris translucentibus, posticarum dimidio apicali lâte cœruleo: subtus dilutioribus; posticis ad marginem internum leviter cœruleo lavatis, lineolis tribus fuscis subparallelis marginatis, fasciis duabus ferrugineis, una ultra cellulas a costa anticarum ad marginem posti-