

1200 feet; and since our return a single example has been sent us from Choctum, in the same district.

We have figured and described the type from the forests of Northern Vera Paz.

g. Wings of male dead blue, of female brown, but wings of both sexes beneath always tinged with blue: first subcostal branch of primaries emitted before the end of the cell, the second just after it: upper discocellular very short; second strongly convex.

28. ***Euptychia glaucina.*** (Tab. VIII. figg. 18, 19.)

Euptychia glaucina, Bates, Ent. Monthl. Mag. i. p. 203¹.

♂ alis supra griseo-cyaneis, venis, costa et margine externo fusciscentibus: posticis lineolis duabus submarginalibus notatis; subtus pallide cyaneis, lineolis duabus submarginalibus et quatuor transversis rufescentifuscis ornatis; linea posticarum tertia ocellos quatuor includente, duos ad angulum apicalem et duos ad angulum analem.

♀ alis supra fuscis; posticis lineolis obsolete marginatis, subtus sicut in mare.

Hab. MEXICO, Jalapa (*Deppe, Mus. Berol.*), Potrero (*Hedemann, Mus. Vindob.*), Valladolid, Yucatan (*Gaumer*); GUATEMALA, Dueñas (*F. D. G. & O. S.*¹), Zapote, San Isidro, and Cahabon (*Champion*).

This is a northern representative of *E. lycidice*, differing, at a glance, from that species in the much greyer tint of the wings of the male. Until quite recently the only specimen known to us was one captured by us at Dueñas in 1861, which served as the type of Mr. Bates's description, and is the one now figured. Within the last few months we have received additional specimens from Yucatan, and also a good supply of examples from Mr. Champion, who captured them chiefly in the neighbourhood of Cahabon, though he also found it elsewhere. It would thus seem to have a considerable range, and in altitude is found from the sea-level to a height of 5000 feet.

h. Secondary wings beneath with a submarginal row of metallic spots or lines; first subcostal branch of primaries emitted before the end of the cell, the second after it.

d. Upper discocellular of primaries short, directed forwards.

29. ***Euptychia philodice.*** (Tab. IX. figg. 15, 16.)

Euptychia philodice, Godm. & Salv. P. Z. S. 1878, p. 264¹.

♂ alis supra fusco-brunneis; posticis macula conspicue irregulari fulva ultra cellulam, ocellis tribus submarginalibus absque pupillis; subtus rubescenti-brunneis, anticis lineis tribus subparallelis ferrugineis, fascia subtriangulari pallide fulva inter duas exteriores marginem internum versus; posticis linea valde sinuata per cellulam, altera extra eandem et extus fascia fulvida late limbata, ocellis duobus submarginalibus ad medium marginis externi, maculas argenteas sagittiformes includentibus, maculis quoque similibus ad marginem reliquum.

♀ mari similis, sed alis paulo dilutioribus et marginibus suis magis rotundatis.

Hab. COSTA RICA, Volcan de Irazu (*Rogers*).