

omnino (aut fere) absentibus; subtus area basali antillarum obscuriore et ocellis intra lineam duplicem postillarum minoribus.

Hab. COSTA RICA (*Van Patten*¹), Irazu (*Rogers*).

Very closely allied to *O. puerta* of Colombia, and *O. simplex* of Colombia and Ecuador; it differs, however, chiefly in the submarginal fulvous band being broader on the secondaries, whilst it becomes very narrow towards the costa of the primaries, which it does not quite reach. Most of our Costa-Rican specimens are either destitute of or only have one or two ill-defined black spots on this band. Beneath also it is very similar to *O. simplex*, but has the inner portion of the primaries blackish brown instead of ferruginous.

The original description was based upon a single specimen in Dr. Van Patten's collection. We have since received several other examples, all agreeing with the type.

7. *Oxeoschistus rogersi*. (Tab. IX. figg. 1, 2.)

Oxeoschistus rogersi, Godm. & Salv. P. Z. S. 1878, p. 267¹.

Alis fuscis, anticis fascia submarginali venis divisa et parte posteriore maculas disjunctas formante, iis inter ramos medianos et radiales superiores fusco notatis, ciliis albescentibus; subtus anticis ut supra, sed dilutioribus et maculis minus definitis, posticis fusco-fulvis, serie ocellorum submarginalium maculisque variis per alam sparsis argenteo-albis.

Hab. COSTA RICA, Irazu (*Rogers*¹); PANAMA, Chiriqui (*Mus. Staudinger*).

The first example of this insect was obtained by our collector Mr. Rogers from the mountains of Irazu, in Costa Rica. Soon after the receipt of this specimen Dr. Staudinger sent us another to name from the neighbourhood of Chiriqui. These are the only ones we have seen at present. Its nearest ally is *O. phaselis* (Hew.), a species placed in the genus *Pronophila* by Hewitson, and subsequently in *Dædalma* by Mr. Butler, but which, in our opinion, ought to be removed to *Oxeoschistus*. From this *O. rogersi* differs at first sight in the tawny markings on the primaries, those wings of *O. phaselis* being of a uniform brown. On the underside the two are very nearly alike.

8. *Oxeoschistus gigas*. (Tab. X. figg. 8, 9.)

Oxeoschistus gigas, Godm. & Salv. P. Z. S. 1877, p. 62¹.

♀ alis fuscis, anticis punctis submarginalibus quinque albis, aliis duobus interioribus marginem internum versus, posticis marginibus late fulvis, venis fuscis divisis, et puncto nigrescente inter ramos medianos primum et secundum notatis; subtus anticis ut supra, sed apice fulvo tincto et maculis indistinctis albidis notatis, posticis fulvescentibus externe pallidioribus, ocellis subapicalibus indistinctis serie positis et maculis variis albidis per alam notatis.

Hab. GUATEMALA, Aceytuno (*Rodriguez*¹), Purula (*Champion*).

The type specimen of this fine species was given to Salvin by Don Juan Rodriguez,