

Hab. GUATEMALA, Polochic valley and Chuacus (*Hague*¹), Senahu, Panima (*Champion*); COSTA RICA (*van Patten*²), Rio Sucio (*Rogers*).

We described this species from examples sent us by Mr. Hague from Guatemala; since then we have likewise received it from Costa Rica. It is very similar to *E. aliphera*, but is rather larger, with the apex of the primaries more rounded, the tawny colour darker, and the black markings much stronger, the secondaries beneath have a row of white spots on the margin, which is not present in *E. aliphera*. As far as we know, it is confined to the localities indicated above.

The specimen figured is one of the types from Guatemala.

5. *Eueides lybioides*.

Eueides lybioides, Staudinger, Verh. z.-b. Ges. Wien, 1875, p. 99¹.

Alis anticis fuscis, fascia arcuata a basi per aream mediam fere ad marginem externum eunte, et macula obliqua apicem propiore fulvis, posticis nigro marginatis.

Hab. PANAMA, Chiriqui (*Ribbé*¹, *Arcé*).

This species was described by Dr. Staudinger, who pointed out how it differed from its very close ally *E. lybia*. These differences, though slight, are constant in all our specimens; and *E. lybioides* appears to constitute a distinct race. On the upperside the subapical tawny spot of the primaries is paler than in *E. lybia*. Beneath, the whole surface is also somewhat paler; and this is especially the case with the subapical spot, which is of a dirty yellow. There is also a faint yellow additional spot at the apex of the primaries; and the black margin of the secondaries is divided by a similarly coloured streak. The base of the costa of the primaries beneath in *E. lybia* is red, but not so in *E. lybioides*.

The range of *E. lybia* extends throughout Guiana and the Amazons basin to Peru, whilst that of *E. lybioides* is confined to the State of Panama.

6. *Eueides olympia*. (Tab. XIX. fig. 1.)

Papilio olympia, Fabr. Ent. Syst. iii. p. 166¹.

Eueides olympia, Butl. Cat. Fabr. Diurn. Lep. p. 120².

Eueides leucomma, Bates, Ent. Monthl. Mag. iii. p. 88³.

E. lybioidi similis, sed macula anticarum prope apicem alba distinguenda.

Hab. NICARAGUA, Chontales (*Belt*, *Janson*); PANAMA, Santa Fé (*Arcé*), Lion-Hill station (*M^cLeannan*³).—W. COLOMBIA.

Mr. Bates described this insect, under the name of *E. leucomma*, from specimens he received from us captured by M^cLeannan on the Panama Railway. There can, however, be no doubt that Fabricius's *Papilio olympia* refers to this species, as Mr. Butler has pointed out²; and therefore the latter name, having priority, must be used. *E. olympia* is very like *E. lybioides*, but may be distinguished from it at a glance by its