

P. atroniæ quoad marem omnino similis.

♀. Feminæ *P. cassiopeæ* et *P. atroniæ* similis, sed anticis fascia ultra cellulam lata alba et posticis fascia alba nulla distinguenda.

Hab. PANAMA, Volcan de Chiriqui (*Ribbe*¹), Bugaba (*Arcé*).

This species too was described by us from a female example in the collection of Dr. Staudinger; and we have since received another from Herr Ribbe, which we now figure. We have also a male, which, as in the case of *P. cassiopea*, does not differ from that sex of *P. atronia*. The three species derive their characters from the females alone, the differences in this sex being associated with a difference of locality.

Regarding *P. chromis* (Godm. & Salv. Ann. & Mag. N. H. ser. 5, ii. p. 260) we are in some doubt, as it now seems to us that it not improbably belongs to *P. diallus*, the chief difference being in the colour of the marks of the primaries. It may, however, represent a fifth species of the *P. atronia* group, having a male like the true *P. atronia*, but a female differing slightly from all the rest.

c. Secondaries above, except outer margin and anal angle, covered with dull sericeous scales.

39. *Phyciodes fulviplaga*. (Tab. XXII. figg. 26, 27.)

Phyciodes fulviplaga, Butl. Cist. Ent. i. p. 77¹; Lep. Ex. p. 183, t. 63. f. 2²; Butl. & Druce, P. Z. S. 1874, p. 348³.

Alis fuscis, anticis macula transversa lata ultra cellulam punctisque marginem versus ochraceis, posticis margine excepto sericeo-fuscis, linea submarginali anguste fulva; subtus area mediana plaga magna fusco-nigra ornata, posticis rufo-fuscis fusco irregulariter marmoratis.

♀. Mari similis, fascia anticarum fulva ad angulum analem producta.

Hab. COSTA RICA (*Van Patten*^{1 2}), Cache (*Rogers*).

A pretty species, with no near allies, which was first discovered by Van Patten and described by Mr. Butler. We have since received many specimens from Mr. Rogers, one of which, a female, we now figure.

40. *Phyciodes otanes*. (Tab. XXII. figg. 28, 29.)

Eresia otanes, Hew. Ex. Butt., *Eresia*, t. 6. f. 47¹.

P. fulviplagæ similis, maculis anticarum fulvis absentibus, colore sericeo posticarum in aream medianam anticarum extendente.

♀ adhuc ignota.

Hab. GUATEMALA (*O. S.*¹), Dueñas and Zapote (*Champion*).

This species was originally described by Hewitson from a specimen obtained in Guatemala by Salvin in 1859, and figured in the 'Exotic Butterflies.'¹ Since then Mr. Champion has obtained several additional specimens at Dueñas and Zapote, the type having most probably been taken at the former place.

We have not yet seen a female of *P. otanes*, which probably resembles that sex of *P. sopolis*. Our figure represents a male from Dueñas.