

*b'*. Apex of primaries elongated, outer margin concave.

## 2. *Eunica caresa*.

*Cybdelis caresa*, Hew. Ex. Butt. *Cybdelis*, iii. f. 20<sup>1</sup>.

*Eunica caresa*, Bates, Journ. Ent. ii. p. 197<sup>2</sup>.

Alis velutino-fusco-nigris, albo extus ciliatis, anticis elongatis apice valde hamatis, area discali et alæ parte basali late purpureo-cyaneis, posticis ad basin eodem colore lavatis; subtus fuscis, anticarum apicibus et posticis omnino rosaceo tinctis et undique albido atomatis, maculis et lineis omnibus valde indistinctis, ocello unico ad anticarum apicem, et posticis quatuor ultra cellulam ornatis, his minutis, parte proxima nigra, parte distali cærulescente.

Femina brunnea, anticis ad apicem obscurioribus fascia lata alba transvittatis, maculis tribus subapicalibus quoque notatis; subtus sicut in mare, maculis paulo clarioribus.

*Hab.* GUATEMALA, Zapote (*Champion*).—COLOMBIA<sup>1</sup>; ECUADOR; PERU; BOLIVIA; AMAZONS<sup>2</sup>.

Colombia is the country whence Hewitson's types were procured; but the range of the species has since been traced as far south as Bolivia and to the valley of the Upper Amazons. We have also recently received from Mr. Champion several examples from the mountains of Guatemala sloping towards the Pacific Ocean; these were taken at an elevation of about 2000 feet above the sea. It is remarkable, however, that no other trace of it has as yet been found in Central America. Our Guatemala specimens have the markings of the secondaries rather more uniform beneath than is usual in South-American examples; but as these latter offer considerable variation in this respect, too much stress must not be laid upon it.

*E. caresa*, though provided with a pencil of hairs near the base of the cell of the secondaries, as in *E. sophronisba*, has the wings of a very different shape, falcate instead of rounded. Its nearest South-American allies are *E. picea* and *E. cabira*, both described from Venezuela by Dr. Felder.

*b*. Male with a pencil of hairs near the base of the submedian nervure of the primaries beneath.

## 3. *Eunica alcmena*. (Tab. XXIII. figg. 6, 7 ♀.)

*Callianira alcmena*, Doubl. & Hew. Gen. Diurn. Lep. t. 28. f. 1<sup>1</sup>.

*Eunica alcmena*, Godm. & Salv. Trans. Ent. Soc. 1880, p. 123<sup>2</sup>.

Alis nigris, fascia subcostali a basin ad angulum apicalem (ad cellulæ finem fere interrupta) cærulea, posticis fascia submarginali eodem colore sed magis micante; subtus fuscescentibus, anticis ad apicem colore pallidiore notatis, posticis rufescenti-fuscis, maculis fusco-nigris variegatis, ocello bipupillato ad angulum apicalem aliisque duobus magnitudine diversis angulum analem propioribus.

Femina alis obscure brunneis, anticis ad apicem obscurioribus fascia transversa ultra cellulam alba.

*Hab.* MEXICO, Cordova (*Rümeli, Höge*); BRITISH HONDURAS, Corosal (*Roe*); GUATEMALA, Polochic valley (*F. D. G. & O. S.*), Teleman (*Champion*), Choctum (*Hague*); PANAMA, Bugaba (*Arcé, Champion*).—COLOMBIA<sup>2</sup>.