

c'. Band of the primaries subparallel to the outer margin.

c''. Transverse lines across the end of the cell of the primaries beneath.

a'''. Band common to both wings fulvous.

8. *Adelpha leuceria*. (Tab. XXX. figg. 7, 8.)

Heterochroa leuceria, Druce, Cist. Ent. i. p. 286¹.

Alis supra brunneis, fascia communi marginibus externis subparallela ad costam anticarum bifida fulva; subtus rufescentibus, fascia sicut supra albicante, serie communi macularum submarginalium indistincte albida; anticis maculis cellularibus duabus et una ultra eam cinereis nigro utrinque marginatis, posticis ad basin fasciis duabus cinereis intima utrinque extima margine interno tantum nigro marginatis.

Hab. GUATEMALA, Polochic valley (*Hague*¹), Sinanja (*Champion*); COSTA RICA, Rio Sucio (*Rogers*).

Mr. Hague supplied us with the types of this pretty species, which were taken in the valley of the Polochic¹; Mr. Champion has since found it in an adjoining valley. It occurs, too, in Costa Rica, where Mr. Rogers captured two specimens.

It has no near allies that we know of. The tawny band running across both wings is a peculiar feature shared only by *A. ethelda* of Ecuador, a species which differs totally from the present one on the underside.

b'''. Secondaries above without conspicuous marks.

9. *Adelpha tracta*.

Heterochroa tracta, Butl. Lep. Ex. p. 102, t. 38. f. 6¹; Butl. & Druce, P. Z. S. 1874, p. 343².

Alis supra brunneis, anticis fascia lata margini externo parallela venis divisa fulva, extra eam fascia altera maculosa indistincta; posticis fasciis tribus margini externo subparallelibus obsolete fulvidis; subtus fere ut in *A. leuceria* fascia posticarum angustiore.

Hab. COSTA RICA (*Van Patten*^{1 2}), Rio Sucio (*Rogers*); PANAMA, Volcan de Chiriqui (*Arcé, Champion*).

The first examples of this insect came in Van Patten's collection from Costa Rica and were described by Mr. Butler¹. We have since obtained many specimens from Mr. Champion taken in Chiriqui, where it appears to be by no means uncommon at an elevation of from 3000 to 4000 feet.

Mr. Butler's plate represents the band of the primaries of far too pale a colour; it is, in fact, tawny brown, a darker tint than is usual in this genus.

Like *A. leuceria*, *A. tracta* is a very isolated species, but they are perhaps more nearly related to each other than to any others in the genus.