

3. *Prepona phædra*, sp. n.

P. demophonti similis, sed posticis ocellis septem submarginalibus notatis; alis subtus multo magis obscuris, fasciola per cellulam anticarum valde sinuata, linea nigra transversa posticarum valde interrupta ad venam subcostalem et ad venam medianam.

Hab. PANAMA, Volcan de Chiriqui (*Trötsch*).

We are indebted to Dr. Staudinger for bringing this species under our notice, as no specimens have as yet been sent to us. It seems certainly distinct from both *P. demophon* and *P. camilla*, the distinct row of submarginal ocelli of the secondaries alone rendering it not difficult of recognition; in addition to this, the transverse black band of the secondaries beneath is almost as much broken as in *P. amphitoe*, which again differs in the silvery tint of the wings beneath.

4. *Prepona amphitoe*.

Nymphalis amphitoe, Godt. Enc. Méth. ix. p. 407¹.

Prepona amphitoe, Bates, Journ. Ent. ii. p. 337².

P. demophonti similis, sed subtus alis argenteo-cinereis (nec ochraceis), linea posticarum transversa interrupta et valde sinuosa dignoscenda.

Hab. GUATEMALA, Dueñas, central valleys (*F. D. G. & O. S.*), San Juan in Vera Paz (*Champion*); NICARAGUA, Chontales (*Belt, Janson*); COSTA RICA (*Van Patten*), Cache (*Rogers*); PANAMA, Chiriqui (*Boucard*), David, Volcan de Chiriqui (*Champion*), Lion Hill (*M'Leannan*).—COLOMBIA to SOUTH BRAZIL.

There is scarcely any difference between this species and *P. demophon* on the upper-side, but beneath it is of a silvery grey (not ochreous), and the lines are always black or dusky, and that crossing the centre of the secondaries, although taking much the same direction as in *P. demophon*, is always strongly angulated, which is not the case in the other species. We find that certain specimens vary much in the same direction as in *P. demophon*, being clouded with an obscure ashy colour.

It is tolerably abundant in Central America; but Mr. Bates says it is less common than *P. demophon* throughout the Amazons region².

♂. Wings beneath divided by a straight black line.

5. *Prepona amphimachus*.

Papilio amphimachus, Fabr. Syst. Ent. p. 457¹.

Morpho amphimache, Hübn. Ex. Schmett. ii. t. 59².

Prepona amphimachus, Bates, Journ. Ent. ii. p. 337³; Butl. & Druce, P. Z. S. 1874, p. 341⁴.

Papilio menander, Cr. Pap. Ex. t. 12. f. A, B⁵.

Prepona menander, Bates, Journ. Ent. ii. p. 336⁶; Butl. & Druce, P. Z. S. 1874, p. 342⁷.

Alis supra *P. demophonti* similibus, subtus ochraceo- aut cinereo-argenteis, linea communi per cellulæ finem anticarum ad angulum analem posticarum nigra, area basali lineis nigris notatis, dimidio apicali fuscescenti.

♀ subtus dimidio apicali pallidiori.