

We have hitherto been unable to find the male of this *Eurygona*, the species apparently being quite a distinct one. The two females originally described are the only ones we have seen.

14. *Eurygona pusilla*. (Tab. XL. figg. 16, 17.)

Eurygona pusilla, R. Felder, Verh. k. k. zool.-bot. Ges. Wien, 1869, p. 467¹.

♀ alis fusco-nigricantibus, anticis area discali flavidо-rufa, posticis fere omnino rufo-brunneis; subtus glauco-albis linea communi, in posticis flexuosa, rufa; anticis triente apicali fusca lunulas albas includente; posticis ad marginem externum punctis nigris notatis, eo ad medium maximo.

Hab. MEXICO, Potrero (*Hedemann*¹); BRITISH HONDURAS, Corosal (*Roe*).

We have a single female specimen from British Honduras, which we have little hesitation in assigning to this species, described from a male obtained by Lieut. Hedemann at Potrero in Southern Mexico. Felder, in his description, compares *E. pusilla* with *E. teleclus*; but the differences are very great, not only from it, but from all other species of *Eurygona* with which we are acquainted.

15. *Eurygona regipennis*. (Tab. XL. fig. 1, 2 ♂, 3, 4 ♀.)

Eurygona regipennis, Butl. & Druce, Cist. Ent. i. p. 103¹; P. Z. S. 1874, p. 352².

Eurygona eurysthenes, Hew. Cat., partim³.

Alis nigris, costa luce purpurea nitentibus, posticis ad marginem internum intense nigris; subtus pallide fuscis, fascia mediana alba, introrsum rufo late limbata, extrorsum rufo-fusco; posticis ocellis ad marginem externum ochraceo cinctis, extus albo limbatis, eo ad medium maximo, et intus purpureo nitente, iis ad angulum analem distortis et elongatis.

♀ alis fuscis, fascia communi lata alba antecarum costam haud attingente ad marginem internum posticarum ducta; subtus mari similis, fascia alba latiori.

Hab. GUATEMALA, Teleman (*Champion*); COSTA RICA (*Van Patten*^{1,2}), Cache (*Rogers*); PANAMA (*Mus. Hewitson*³).

Of this beautiful species we have the type, a male in rather poor condition and a female, both from Costa Rica, and also another female from Guatemala, the only one obtained by Mr. Champion.

The female on the upper side recalls *E. eubages*, Hew., but the markings of the under surface show that it has no near relationship with that species.

In the neighbourhood of Chiriqui its place is perhaps taken by *E. eupepla*, the females being very similar. The males, however, differ essentially on the underside.

A female from Panama in the Hewitson collection stands under the name of *E. eurysthenes*³.

16. *Eurygona eupepla*, sp. n. (Tab. XLIII. figg. 5, 6 ♂, 7, 8 ♀.)

Alis supra iis *E. regipennis* similibus, lætissime purpureis, posticis plerumque nigerrimis margine externo tantum purpureo; subtus ochraceo-fulvis, fascia communi ultracellulari rufa in posticis refracta; anticis fascia

Biol. Centr.-Am., Rhopal., Vol. I., December 1885.