

*b.* Sexes dissimilar.*a'*. Anal angle of secondaries rounded.**6. *Lemonias pelarge*.** (Tab. XLIV. figg. 17, 18.)*Lemonias pelarge*, Godm. & Salv. P. Z. S. 1878, p. 366<sup>1</sup>.

Anticis ad basin et posticarum margine externo fusco-nigris, illis maculis indistinctis apicem versus cæruleis, macula ad angulum analem et harum area interna (basi excepta) rufo-aurantiis; subtus griseo-cæruleis fusco-nigro maculatis, margine externo anticarum late, posticarum anguste, fusco-nigris, abdomine supra nigricante, infra griseo.

♀ adhuc ignota.

*Hab.* GUATEMALA, Polochic valley (*Hague*<sup>1</sup>), Cahabon (*Champion*).

This species is closely allied to *L. parthaon* of Guiana and the Amazons valley; the blue spots of the wings are smaller, there is an orange-red spot near the anal angle of the primaries, and this colour extends almost to the very base of the secondaries. Beneath, the primaries have more blue-grey, and the submarginal spots of the secondaries are obsolete; moreover, the abdomen above is black instead of being rufous, as in the allied species.

Our first example of this *Lemonias* was sent us from the Polochic valley by Mr. Hague, a second specimen was taken by Mr. Champion near Cahabon in the adjoining valley. These are all we have seen.

**7. *Lemonias cilissa*.***Lemonias cilissa*, Hew. Ex. Butt., *Lemonias*, 4. f. 33, 34<sup>1</sup>; Bates, Journ. Linn. Soc. Zool. ix. p. 447<sup>2</sup>.

Alis rufo-fuscis, anticis punctis tribus subapicalibus albis, area mediana ad marginem internum et posticis fulvis, lineolis nigris transversim notatis; subtus nigro maculatis, maculis submarginalibus albis, anticis fuscis ad marginem internum pallide ochraceis, posticis (præter marginem externum fuscum) albis.

♀ alis pallide cervinis maculis transversis tenuissimis vix notatis, anticis margine externo late fusco, maculas submarginales albas includente, posticis maculis albis fusco cinctis in serie submarginali positis; subtus fere ut supra sed paullo pallidioribus.

*Hab.* NICARAGUA<sup>2</sup>, Chontales (*Belt*<sup>1</sup>); PANAMA, Chiriqui (*Ribbe*), Bugaba (*Champion*).

Except in Costa Rica, where the species does not appear to exist, *L. cilissa* is very common from Nicaragua to the district of Chiriqui in the State of Panama. We have received a large number of specimens from the latter locality.

Some variation is noticeable amongst them, the secondaries of the males, especially in Nicaraguan specimens, are darker at the base; intermediate examples occur everywhere. Hewitson's types are from Nicaragua.

Frequents low bushes in the second-growth woods of the "terra caliente" (*Champion*).

**8. *Lemonias pseudocrispus*.**

*Lemonias pseudo-crispus*, Westw. in Doubl. & Hew. Gen. Diurn. Lep. p. 459<sup>1</sup>; Butl. Journ. Linn. Soc. Zool. ix. p. 216, t. 6. ff. 9, 10<sup>2</sup>; Bates, Journ. Linn. Soc. Zool. ix. p. 447<sup>3</sup>.