

*b''*. Wings grey beneath.

### 6. *Theope thootes*.

*Theope thootes*, Hew. Exot. Butt., *Theope*, ff. 9, 10<sup>1</sup>; Bates, Journ. Linn. Soc. Zool. ix. p. 454<sup>2</sup>.

Anticis nigricantibus, cellula et infra eam ad marginem internum, posticisque omnino, læte cæruleis, subtus fuscescentibus.

♀ mari similis sed colore anticarum cyaneo magis extenso et extus bifurcato, subtus anticis paullo saturatoribus.

*Hab.* NICARAGUA, Chontales (*Belt*).—AMAZONS VALLEY<sup>12</sup>.

We have a single male specimen from Nicaragua which agrees with our series from the Amazons, the country of the true *T. thootes*. The anal angle of the secondaries is rounded, and there are no spots in this region. These characters separate *T. thootes* from *T. theutis*, a species we have long confounded with it.

*b*. Wings beneath with spots near the anal angle of secondaries, or crossed with a common dark band.

*c'*. Secondaries beneath with spots near the anal angle.

### 7. *Theope theutis*, sp. nov. (*T. thootes*, Tab. XLVI. figg. 16, 17 \*.)

*T. thooti* similis sed alis subtus paullo saturatoribus, posticis angulo anali magis producto haud rotundato maculisque minutis interdum fere obsoletis notato.

*Hab.* GUATEMALA, Polochic valley (*Hague*); NICARAGUA, Chontales (*Belt*).

We have a single female specimen of this species from Guatemala, and a male and two females from Nicaragua. At first sight it may readily be mistaken for *T. thootes*, but the points given above fairly distinguish it.

### 8. *Theope folia*, sp. n. (Tab. XLVII. figg. 23, 24.)

Precedenti similis, sed anticis magis acutis, costa magis sinuosa, colore cyaneo magis nitente; subtus cinereo-fuscis, extus pallidioribus.

♀ nobis ignota.

*Hab.* PANAMA, Peña Blanca (*Champion*).

There is a specimen in the British Museum, said to be from Rio Janeiro, but the name of the collector has not been recorded. Our type is the only one captured by Mr. Champion during his recent journey.

### 9. *Theope herta*, sp. n. (Tab. XLVII. figg. 25, 26.)

Alis fuscis ad basin cæruleo atomatis; subtus sericeo-murinis, maculis parvis ad angulum posticarum analem.

*Hab.* PANAMA, Bugaba (*Champion*).

This species rests upon a single example obtained by Mr. Champion. It has no near

\* These figures are erroneously marked on the Plate as 16 ♂, 17 ♀; they refer to the upper and under side of the same insect, a female of *T. theutis*.