

2. **Opharus albipunctatus**, sp. n. (Tab. X. fig. 11.)

Primaries uniform pale drab brown; secondaries whitish hyaline, with the outer and inner margins bordered with the same colour as the primaries; a very minute blue spot at the base of the primaries: head, palpi, antennæ, and thorax black; the collar spotted with bright blue spots; the patagia yellowish white, with a minute bright blue spot at the extreme point; abdomen yellow, banded with black, also a narrow central black line from the anus to the base, where the abdomen is dusky brown; the underside brown; legs black. Expanse $1\frac{3}{4}$ inch.

Hab. BRITISH HONDURAS, Belize (*Blancaneaux*); GUATEMALA, Panzos, Las Mercedes 3000 feet (*Champion*).

This pretty species is very distinct from any other with which I am acquainted.

3. **Opharus mundator**, sp. n. (Tab. X. fig. 10.)

Primaries uniform semitransparent black; the secondaries similar, but slightly whitish near the base: head, palpi, and antennæ black; thorax black, with two white spots in front, just behind the head; abdomen black, banded with yellowish lines on each segment; a row of white spots on each side; legs black. Expanse $2\frac{1}{2}$ inches.

Hab. COSTA RICA, Irazu 6000 to 7000 feet (*Rogers*).

Only a single example of this species was sent.

4. **Opharus ruficollis**, sp. n. (Tab. X. fig. 8.)

Primaries uniform pale brown, with an indistinct broken darker brown band from the costal margin near the apex to close to the anal angle, the veins all darker; secondaries pale brown, pinkish yellow from the base to beyond the middle: head and front of thorax bright orange-red, with several very minute black spots; thorax brown; abdomen pinkish yellow, spotted with black on the sides; legs brown. Expanse 2 inches.

Hab. PANAMA, Volcan de Chiriqui (*Ribbe, mus. Staudinger*).

I have only seen a single example of this insect.

ANAXITA.

Anaxita, Walker, Cat. iii. p. 748.

Two species have been placed in this genus.

1. **Anaxita decorata**.

Anaxita decorata, Walk. Cat. iii. p. 748¹; Herr.-Schäff. Samml. aussereur. Schmett. f. 528.

Hab. MEXICO (*Hartweg*)¹.

The only specimens of this very fine species known to me are in the British Museum; it is well figured by Herrich-Schäffer.

Subfam. *PERICOPIINÆ*.

ESTHEMA.

Esthema, Hübner, Verz. bek. Schmett. p. 178.

This genus is represented in our country by three species, two of them being common South-American forms.