

12. *Letis hypnois*.

Syrnia hypnois, Hübn. Samml. exot. Schmett. ii. t. 210. ff. 3, 4; Verz. bek. Schmett. p. 273; Guén. Sp. gén. des Léop. vii. p. 157¹; Walk. Cat. xiv. p. 1277².

Hab. MEXICO, Coatepec (*Brooks*), Jalapa (*Schaus*); BRITISH HONDURAS, Belize, Cayo (*Blancaneaux*); GUATEMALA, Las Mercedes (*Champion*), Coban (*Conradt*); PANAMA, Chiriqui (*Ribbe, in mus. Staudinger*).—COLOMBIA^{1 2}; AMAZONS, Pará²; BRAZIL^{1 2}, Santa Catarina.

From Central America we have only received a few specimens of this fine species. I have not the least doubt that it is a *Letis*.

LATEBRARIA.

Latebraria, Guénée, Sp. gén. des Léop. vii. p. 159 (1852); Walker, Cat. xiv. p. 1281.

In Central America this genus is represented by one species. Walker, in his Catalogue, included six in it; at least three of these belong to other genera or to previously known species. The two species described by Guénée are from Tropical America or the Antilles.

1. *Latebraria amphipyroides*.

Latebraria amphipyroides, Guén. Sp. gén. des Léop. vii. p. 159, Noct. t. 17. f. 2¹; Walk. Cat. xiv. p. 1282².

Latebraria errans, Walk. Cat. xiv. p. 1281³.

Hab. MEXICO, Jalapa (*Schaus*), Coatepec (*Brooks*), Omilteme in Guerrero 8000 feet (*H. H. Smith*), Yucatan³, Valladolid in Yucatan (*Gaumer*); GUATEMALA, Las Mercedes 3000 feet, Panzos in Vera Paz (*Champion*); COSTA RICA, Volcan de Irazu 6000 to 7000 feet (*Rogers*).—COLOMBIA^{1 2}; VENEZUELA³; ECUADOR; BRAZIL^{1 2}; WEST COAST OF AMERICA³; ANTILLES, Cuba^{1 2}.

This species varies considerably in colour, some examples being much paler than others; in our region it is a fairly common insect. Mr. Smith met with it in the month of August.

CYCLOPIS.

Cyclopis, Hübner, Verz. bek. Schmett. p. 274 (1816); Guénée, Sp. gén. des Léop. vii. p. 165; Walker, Cat. xiv. p. 1288.

The single species of this genus found in our region appears to be very common in Guatemala, but from the State of Panama we have as yet only received one specimen.

1. *Cyclopis cæcutiens*.

Cyclopis cæcutiens, Hübn. Samml. exot. Schmett. ii. t. 213. ff. 3, 4; Verz. bek. Schmett. p. 274; Guén. Sp. gén. des Léop. vii. p. 166¹; Walk. Cat. xiv. p. 1288²; Möschl. Verh. zool.-bot. Ges. Wien, 1880, p. 429³; Beitr. zur Schmett.-Fauna von Surinam, iii. p. 51⁴.