

### 5. *Conchylodes (?) sabatalis*, sp. n. (Tab. LXII. fig. 18.)

Primaries and secondaries pale yellowish-white; the primaries crossed from the costal to the inner margin by four dark brown lines, which are broken into spots, the outer margin from the apex to the anal angle shaded with dark brown, the fringe pale yellowish-brown; the secondaries crossed by two rows of broken brown lines, the apex slightly edged with brown, the marginal line black, the fringe pale yellow: head, thorax, and abdomen yellow, the anus black, the antennæ and legs yellowish-white.—*Female*. Very similar to the male. Expanse 1 inch.

*Hab.* MEXICO, Atoyac in Vera Cruz, Dos Arroyos in Guerrero (*H. H. Smith*), Temax in North Yucatan (*Gaumer*); GUATEMALA, El Reposo (*Champion*).

The Guatemalan specimens are much smaller than those from Mexico. An example from Dos Arroyos is figured.

### 6. *Conchylodes levinia*.

*Phalaena Pyralis levinia*, Cram. Pap. Exot. iv. p. 131, t. 357. fig. K<sup>1</sup>.

*Spilomela levinialis*, Guen. Sp. gén. des Lép. viii. p. 283<sup>2</sup>.

*Zebronia levinialis*, Walk. Cat. xvii. p. 472<sup>3</sup>.

*Conchylodes levinia*, Leder. Wien. ent. Monats. vii. p. 443<sup>4</sup>.

*Zebronia bunusalis*, Walk. Cat. xix. p. 967<sup>5</sup>.

*Conchylodes bunusalis*, Warr. Trans. Ent. Soc. Lond. 1889, p. 267<sup>6</sup>.

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith*); PANAMA, Chiriqui (*Ribbe, in mus. Staudinger*).—ECUADOR; GUIANA, Surinam<sup>3 4</sup>, Cayenne<sup>2 4</sup>; AMAZONS<sup>6</sup>; BRAZIL, Rio Janeiro<sup>5</sup>.

We have received three specimens of this species from Mexico. An example from Ecuador in my own collection has the wings more heavily spotted.

### COENOSTOLA.

*Cœnosta*la, Lederer, Wien. ent. Monats. vii. p. 408 (1863).

#### 1. *Cœnosta*la apicalis.

*Cœnosta*la apicalis, Leder. Wien. ent. Monats. vii. pp. 409, 479, t. 14. fig. 16<sup>1</sup>; Walk. Cat. xxxiv. p. 1484<sup>2</sup>.

*Hab.* PANAMA, Chiriqui (*Ribbe*), Volcan de Chiriqui 2000 to 3000 feet (*Champion*).—COLOMBIA; BRAZIL<sup>1 2</sup>.

Mr. Champion captured a few specimens of this insect on the Volcan de Chiriqui. Examples from Bogota in my own collection are slightly larger.

### OMIODES.

*Omiodes*, Guenée, Sp. gén. des Lép. viii. p. 355 (1854); Lederer, Wien. ent. Monats. vii. p. 409.

A very distinct and well-marked genus; the sexes differ considerably in some of the species.