

1. *Leucochroma splendidalis*.

Phalena Pyralis splendidalis, Cram. Pap. Exot. iv. p. 161, t. 371. fig. L¹.

Epipagis splendidalis, Hübn. Verz. bek. Schmett. p. 358².

Leucochroma splendidalis, Guen. Sp. gén. des Lép. viii. p. 287³; Walk. Cat. xvii. p. 491⁴.

Botys splendidalis, Leder. Wien. ent. Monats. vii. p. 371⁵.

Botys selectalis, Walk. Cat. xxxiv. p. 1396⁶.

Leucochroma minoralis, Warr. Trans. Ent. Soc. Lond. 1889, p. 267⁷.

Hab. MEXICO, Paso de San Juan in Vera Cruz (*Schaus*); HONDURAS⁴, Ruatan I. (*Gaumer*); PANAMA, Volcan de Chiriqui 2000 to 3000 feet (*Champion*).—COLOMBIA, Bogota⁶; ECUADOR; GUIANA, Surinam³, Cayenne³; AMAZONS⁷; ANTILLES, San Domingo⁴.

In some specimens of this species the yellow markings on the wings are very faint.

2. *Leucochroma prosalis*, sp. n. (Tab. LXII. fig. 30.)

Primaries and secondaries yellowish-white, the primaries with the base and the inner margin spotted with yellowish-brown, a large yellowish-brown patch extending across the wing near the apex almost to the anal angle, on the outer edge of which is a pale waved line, the marginal line spotted with black; the secondaries with several dark brown spots partly crossing the wing from the anal angle; the fringe yellowish: head, antennæ, thorax, abdomen, and legs yellowish-white. Expanse 1 inch.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).

One specimen.

3. *Leucochroma saltigalis*, sp. n. (Tab. LXII. fig. 31.)

Primaries and secondaries silky-white; the primaries spotted with yellow at the base and in the cell, the outer margin broadly bordered with yellowish-brown, the border paler near the apex and preceded by a waved dark brown line, which crosses the wing from the costal to the inner margin, the marginal line spotted with black at the apex, the fringe pale brown; the secondaries crossed about the middle by a row of yellow spots, and with a submarginal waved brown line extending partly across the wing from near the apex, but not reaching the anal angle, a small black spot on the anal angle, and a black marginal line at the apex, the fringe white, yellowish at the apex: head, antennæ, abdomen, and legs white; the thorax white, spotted with yellow. Expanse $1\frac{1}{4}$ inch.

Hab. PANAMA, Chiriqui (*Trötsch, in mus. Staudinger*).

One specimen of this very distinct and beautiful species is contained in Dr. Staudinger's collection. It is allied to *L. splendidalis* (Cr.).

4. *Leucochroma (?) ruscialis*, sp. n. (Tab. LXIII. fig. 1.)

Primaries and secondaries semihyaline white; the primaries with a spot on the costal margin and one on the inner margin, both close to the base, two lines in the cell and one below it, a large spot near the anal angle, and a spot on the outer margin, black, a curved black line extending from the costal margin to the anal angle; the secondaries with a small black streak on the middle of the outer margin and a small black dot on the inner margin above the anal angle: head, antennæ, thorax, abdomen, and legs white. Expanse $1\frac{3}{10}$ inch.

Hab. PANAMA (*Boucard, in mus. D.*), Chiriqui (*Arcé, in mus. D.*).

This species is quite unlike any other known to me.