

Syntomedia epilais (I. p. 43).

To the localities given, add:—MEXICO, Jalapa (*M. Trujillo*), Orizaba (*Boucard, in mus. D.*); BRITISH HONDURAS, Belize (*Sir Alfred Moloney, in mus. D.*), Corosal (*Roe, in mus. D.*).

5. Syntomedia vulcana. (Tab. LXX. fig. 25.)

Syntomedia vulcana, Druce, Ann. & Mag. Nat. Hist. (6) iv. p. 83 (1889) ¹.

Primaries uniformly deep glossy black, with a rich purplish tinge in some lights; a small streak in the cell and a round spot near the base on the inner margin very pale primrose-colour; a white dot close to the base on the costal margin; secondaries deep glossy black, the basal part hyaline, as in *S. melanthus* (Cram.): head, thorax, and abdomen glossy bluish-black, the abdomen with a row of white spots on each side; the legs and antennæ black, the antennæ with the tips white on the underside. Expanse $2\frac{2}{10}$ inches.

Hab. MEXICO, Tierra Colorada in Guerrero 2000 feet (*H. H. Smith*).

One specimen, captured in October 1888. Allied to *S. melanthus* (Cram.), and also to *S. sauleyi*, Guen., but very distinct from either.

6. Syntomedia joda, sp. n. (Tab. LXXI. fig. 15.)

Primaries and secondaries deep black; the primaries with a round spot about the middle and a wide band partly crossing the wing beyond it chrome-yellow; the secondaries with the base and a round spot about the middle chrome-yellow; the fringe black; the underside of both wings as above: head, thorax, legs, and antennæ black, the antennæ tipped with white, the collar, tegulæ, and base of the thorax chrome-yellow; abdomen above glossy blue-black, with two of the segments banded with chrome-yellow, beneath dull black, with one segment only banded with yellow. Expanse $1\frac{1}{2}$ inch.

Hab. MEXICO, Guadalajara (*coll. Schaus*).

A very distinct species, resembling in some respects the African *Euchromiæ*.

ENDERA (I. p. 43).

Endera vulcanus (I. p. 43).

To the locality Mexico, add:—Cuesta de Misantla (*M. Trujillo*), Amula in Guerrero 6000 feet (*H. H. Smith*).

A rare species, of which we have received very few specimens.

EUPYRA (I. p. 43).

Eupyra principalis (I. p. 43). (Tab. LXX. fig. 27, ♂.)

To the locality given, add:—MEXICO, Sierra Nevada de Colima 12,000 feet, Lake Chapala in Jalisco (*Richardson*), Jalisco (*Schumann*), Amecameca in Morelos (*F. D. G.*), Amula in Guerrero 6000 feet (*H. H. Smith*); GUATEMALA, in the city (*Rodriguez*).

Varies considerably in colour, some specimens being much brighter than others. We figure a male from Jalisco.

Eupyra corydon (I. p. 44).

To the locality given, add:—MEXICO, Jalapa (*M. Trujillo*), Orizaba (*Flohr; Boucard, in mus. D.*).