

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith* <sup>1</sup>).

Allied to *D. mexicana*, but differing from it in the coloration of the thorax and abdomen.

1 (B). **Dycladia thera.** (Tab. LXXI. fig. 26.)

*Dycladia thera*, Druce, Ann. & Mag. Nat. Hist. (6) iv. p. 85 (1889) <sup>1</sup>.

Primaries black, with the hyaline markings almost identical with those of *D. mexicana*, but with the apical spot smaller and narrower; the secondaries with the black margin considerably narrower than in that species; the underside of both wings as above: head and collar bright blue, the thorax and upperside of the abdomen dull black; the tegulæ black, with a yellow dot at the base; the abdomen with a yellow spot on each side close to the base, the sides and the anal segment bright blue, the underside and the anal tuft bright scarlet; antennæ and legs black. Expanse  $1\frac{1}{2}$  inch.

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith* <sup>1</sup>).

Allied to *D. mexicana*, but larger, and differing in the coloration of the head, thorax, and abdomen.

1 (c). **Dycladia demona**, sp. n. (Tab. LXXI. fig. 29.)

Primaries deep black, darkest close to the base, the fringe black; secondaries hyaline, with the veins, apex, and outer margin black: head bright metallic-blue; antennæ black, banded with white near the tip; collar, tegulæ, thorax, and base of the abdomen chrome-yellow, the rest of the upperside of the abdomen metallic blue-black, with a row of orange spots on each side of the three middle segments; underside of the thorax, and the coxæ and femora of the fore legs, bright carmine; underside of the abdomen black. Expanse  $1\frac{4}{10}$  inch.

*Hab.* MEXICO, Jalapa (*coll. Schaus*).

Allied to *D. lydia*, Druce, but very distinct therefrom.

**Dycladia vittata** (I. p. 61).

To the localities given, add:—MEXICO, Rinconada and Orizaba in Vera Cruz (*coll. Schaus*), Temax in North Yucatan (*Gaumer*).

**Dycladia correbioides** (I. p. 61).

To the localities given, add:—MEXICO, Jalapa (*M. Trujillo*), Orizaba (*Boucard, in mus. D.*), Misantla (*F. D. G.*), Coatepec (*Brooks*), Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*); GUATEMALA, in the city (*Rodriguez*); HONDURAS (*Candèze, in mus. D.*); COSTA RICA, Santa Clara Valley 1200 feet (*Zurcher*), Candelaria Mts. (*Underwood*).

The specimen from Honduras is much more hyaline than any of the others I have seen, and it may prove to belong to a distinct species.

**Dycladia imitata** (I. p. 62).

To the locality given, add:—HONDURAS (*coll. Candèze*).

**Dycladia augusta** (I. p. 62).

To the locality given, add:—MEXICO, Jalapa (*M. Trujillo*).