

the middle, which does not reach the anal segment, the anus black, the underside bright yellow.—*Female*. Primaries dark brown; secondaries pinkish-red, broadly bordered with brown from the apex to the anal angle: head, thorax, and abdomen as in the male. Expanse, ♂ $2\frac{3}{4}$, ♀ $3\frac{1}{4}$ inches.

Hab. MEXICO, Temax in North Yucatan (*Gaumer*¹).

9 (A). **Pericopis humeralis.**

Pericopis humeralis, Walk. Cat. ii. p. 348 (1854)¹.

Hab. MEXICO, Las Vigas (*coll. Schaus*).

Walker's type was from an unknown locality¹.

9 (B). **Pericopis montezuma.** (Tab. LXXVII. figg. 25, 26, ♀.)

Pericopis montezuma, Schaus, P. Z. S. 1892, p. 283 (♀)¹.

Hab. MEXICO, Las Vigas (*Schaus*¹).

Probably a variety of *P. humeralis*, Walk.

Pericopis thetis (I. p. 112).

To the localities given, add:—MEXICO, Cuernavaca in Morelos (*H. H. Smith*), Jalapa (*M. Trujillo*), Orizaba (*Boucard, in mus. D.*).

According to Mr. H. Edwards (Ent. Amer. ii. p. 165), this insect is quite common in some parts of Mexico.

10 (A). **Pericopis howardi.** (Tab. LXXVII. fig. 24, ♂.)

Daritis thetis, Klug, var. *howardi*, H. Edwards, Ent. Amer. ii. p. 165 (1886)¹.

Hab. NORTH AMERICA, New Mexico¹.—MEXICO, Durango city (*Becker*), Guadalajara in Jalisco (*Goldsmith, in mus. D.*).

Altogether a larger and finer insect than *P. thetis* (Klug). A specimen from Durango city is figured.

Pericopis fenestrata (I. p. 112).

To the locality given, add:—GUATEMALA, in the city (*Rodriguez*); HONDURAS (*Wittkugel, in mus. Staudinger*); COSTA RICA, Candelaria Mts. (*Underwood*).

Pericopis umbra (I. p. 113).

To the locality given, add:—HONDURAS, San Pedro Sula (*Wittkugel, in mus. D.*).

Pericopis angulosa (I. p. 113).

To the localities given, add:—MEXICO, Orizaba (*coll. Schaus*); HONDURAS (*Wittkugel, in mus. Staudinger*).