

9. *Polyortha tersa*, sp. n. (Tab. VIII. fig. 14.)

*Antennae* tawny grey, spotted above with fuscous. *Palpi* hirsute above and beneath, extending more than the length of the head beyond it; dark tawny, with some paler speckling at the sides; terminal joint naked, tipped with whitish. *Head* dark tawny, with paler speckling. *Thorax* dark tawny; tufted behind. *Forewings* rather narrow in proportion to their length, termen slightly sinuate below the apex; dark tawny, with a white mediocostal patch, in which is a tawny shade and some tawny scales on the margin—this patch is semiovate and extends to the lower edge of the cell; the dark portion of the wing shows about five strong tufts of raised scales, of which the most conspicuous are on the fold and above the commencement of the dorsal cilia; cilia tawny, with slight purplish grey sheen. *Exp. al.* 16 mm. *Hindwings* greyish fuscous; cilia paler. *Abdomen* ochreous. *Legs* shining, white, with tawny tarsal bands.

*Type* ♂ (66278) Mus. Wlsm. (Godm-Salv. Coll.) BM.

*Hab.* GUATEMALA: QUEZALTENANGO: Cerro Zunil, 4000–5000 ft., VIII–IX. 1880 (G. C. Champion). Unique.

10. *Polyortha euchlorana*, sp. n.

*Antennae* pale ochreous, dotted with fuscous. *Palpi* projecting the length of the head beyond it, median joint hirsute, terminal joint naked; bright green. *Head* and *Thorax* bright green; the latter tufted behind. *Forewings* slightly sinuate below the depressed apex, somewhat narrower than is usual in this genus; bright green, with a broad, mottled, fuscous band crossing the wing at two-thirds, obliquely parallel with the termen; this band contains some tawny grey shading on its dorsal, and some black streaklets on its costal half—two small tawny fuscous costal spots lie between it and the base, some black scales lying also on the costa between them; the wing-surface exhibits several patches of strongly raised scales, at least three on the green basal half, and two on the darker transverse band, conforming in each case to the colour on which they are set; cilia bright green above and beneath; underside dusky, with the exception of some ochreous speckling along the costa and their bright green cilia. *Exp. al.* 12 mm. *Hindwings* 1; tawny grey; with a pale slender line at the base of the grey cilia; underside dusky. *Abdomen* tawny grey. *Legs* pale cinereous.

*Type* ♀ (66280) Mus. Wlsm. (Godm-Salv. Coll.) BM. [PT. (66281) US. Nat. Mus.]

*Hab.* MEXICO: TABASCO: Teapa III. (H. H. Smith). Two specimens.

A very small and beautiful species—quite distinct.

11. *Polyortha dolosa*, sp. n.

*Antennae* fawn-brown. *Palpi* extending scarcely the length of the head beyond it, the median joint thickly clothed beneath, the terminal short obtuse; fawn-brown. *Head* and *Thorax* fawn-brown. *Forewings* with the costa abruptly arched at the base, ciliate from the humeral angle outward; fawn-brown, with a slight rosy tinge, a series of small black marginal spots along the rounded apex and convex termen, a blackish spot on the cell a little beyond its middle; there are also one or two black scales beyond the end of the cell and on the costa, with a thin and scattered sprinkling of bright chestnut scales, especially about the tornus; cilia concolorous with the wings. *Exp. al.* 14 mm. *Hindwings* brownish fuscous; cilia pale cinereous, brownish grey on their basal half. *Abdomen* rather shining brownish. *Legs* shining, pale fawn.

*Type* ♂ (66283, San Gerónimo) Mus. Wlsm. (Godm-Salv. Coll.) BM. [PT. (66284) US. Nat. Mus.]

*Hab.* AMERICA, C-S. Central America—GUATEMALA: BAJA VERA PAZ: San Gerónimo, 1879–90 (G. C. Champion). South America—COLOMBIA: CUNDINAMARCA: Bogota (Birchall). Three specimens.

Very similar to *Tortrix apatela* Wlsm. (p. 289), but easily separable by the neuration.