

RHYPAROCHROMUS (p. 215).

Rhyparochromus plenus (p. 216).

To the locality given, add :—GUATEMALA, Quiché Mts. 8000 feet (*Champion*).

PETISSIUS, gen. nov. (To follow the genus *Trapezus*, p. 217.)

Allied to *Trapezonotus* and *Trapezus*; from the first it differs by having the pronotum much broader than long and the head longer than broad, and from *Trapezus* by having the head, including the outer margins of eyes, about equal in width to the anterior margin of the pronotum. From both and from all the allied genera it is distinguished by the shape of the pronotum, which is much narrower at apex than at base, with the lateral margins non-ampliated. The first and second joints of the antennæ are sparingly hirsute; the anterior femora are spinous beneath at apex, and the intermediate and posterior tibiæ are longly setose.

1. **Petissius assimilandus**, n. sp. (Tab. XXXV. fig. 21.)

Head black; antennæ ochraceous, with the apical joint infuscated, second joint slightly longer than the third, which is longer than the fourth. Pronotum ochraceous, with the anterior lobe mostly black and impunctate, the disk coarsely and darkly punctate, the punctures forming some longitudinal fasciæ on the posterior lobe, and being particularly thick on the sublateral and subanterior margins. Scutellum brownish, thickly covered with coarse black punctures, and with a central levigate and two lateral ochraceous fasciæ. Corium brownish, thickly covered with coarse black punctures, margins of clavus, margins of corium, and the veins ochraceous, and with a few small discal ochraceous spots. Membrane dark fuscous. Body beneath blackish; legs and rostrum ochraceous.

Long. 7 millim.

Hab. GUATEMALA, Mirandilla (*Champion*); PANAMA, Bugaba (*Champion*).

We have received four examples of this species.

2. **Petissius (?) diversus**, n. sp. (Tab. XXXV. fig. 22.)

Body above reddish-ochraceous and pilose; eyes black; scutellum and apical areas of the corium castaneous; antennæ with the basal joint reddish-ochraceous, the remaining joints dull brownish; second joint a little longer than the third, third and fourth joints subequal in length; membrane pale ochraceous; body beneath reddish-ochraceous, legs pale ochraceous. The lateral margins of the pronotum and corium are very narrowly pale ochraceous; the pronotum has a transverse impression; the corium is strongly pilose.

Long. 3 millim.

Hab. GUATEMALA, Paraiso, El Reposo (*Champion*); PANAMA, David (*Champion*).

We have received three (carded) specimens of this species, for which I hesitate to propose a new genus. It differs from the type of *Petissius* by having the head including the eyes a little narrower than the anterior margin of the pronotum; the other characters are, however, generally the same.

EREMOCORIS (p. 218).

2. **Eremocoris germanus**, n. sp. (Tab. XXXV. fig. 24.)

Head black; pronotum with the anterior lobe castaneous, with its lateral margins ochraceous, the posterior lobe black, with an ochraceous spot near the lateral angles. Scutellum pale castaneous, with the lateral