

Simple, pinnate, pinnules simple, elegantly incurved; axis of the stem nearly triangular, of pinnules setaceous.

Coast of Martinique. *Lamarck.* *Lamouroux.*

Antipathes eupteridea, Lamarck, ii. 483, No. 19. une belle plume de paon décolorée et brunnâtre."

—, Lamour. Encycl. 71; "Resemble à

III. *In plano ramosæ.*

1. *Ramis pinnatis.*

5. *ANTIPATHES PECTINATA.* (*Lamarck.*)

A. flabellata, 9" alta; *ramis pinnato-pectinatis*; *ramorum axe compresso*, *ramulorum filiformi*, *subulato*; *spinulis paucis*.

Flabellate, 9 inches high; branches pinnato-pectinate, axis of the branches compressed; of the branchlets filiform, subulate; spinules few.

Antipathes pectinata, Lamk. ii. 480, No. —, Lamour. Pol. flex. 381; Encyc. 6.—Locality not given. 72.

6. *ANTIPATHES MYRIOPHYLLA.* (*Pallas.*)

A. grandis (18–20" alta), latè ramosissima, incurvata, *in plano paniculata*, *subtripinnata*, *pinnulis brevibus* (vix $\frac{3}{4}$ "), *crebris*; *pinnularum axe setaceo*, *scabro*.

Large (18 to 20 inches high), much and spreading rameous, incurvate, paniculate in a plane, subtripinnate, pinnules short ($\frac{3}{4}$ of an inch or less), much crowded; axis of the pinnules setaceous, scabrous.

Indian Ocean. *Ellis.* *Lamouroux.*

- Myriophyllum Indicum ramosissimum*, *Antipathes myriophylla*, Esper, Fortsetz. Pettiv. Gazoph. tab. 35, fig. 12. i. tab. 10.
Antipathes myriophylla, Pallas, Zooph. 210. —, Lamk. ii. 481, No. 9.
 —, Ellis and Sol. 102, tab. 19, figs. 11, 12.—From the Moluccas. —, Lamour. Pol. flex. 378; Exp. 32, pl. 19, figs. 11, 12; Encyc. 71. —, Blainv. Man. 510, pl. 87, fig. 2.