

*plumæ, subtùs pallidiores ; scapis supernè nigris, subtùs albis : rostro unguibusque cærulescenti-nigris : cerâ, palpebris, tibiis cambogiaconitidis : iride aurantiâ. Mas adultus.*

*A. juvenis : capite dorsoque umbrinis, plumis ferruginoso marginatis, albo fasciatis, fasciis vix discernendis nisi plumæ erigantur : gutture albo : pectore albidescenti-fulvo, singulas plumas lineâ medianâ maculâque apicali umbrinâ notantibus : lateribus umbrino fasciatis ; femoribus fulvis, pallido ferrugineo irregulariter fasciatis : alis umbrinis, fasciis saturatioribus ; subtùs fulvis, umbrino vittatis ; scapis subtùs furvis, supernè brunneis : caudâ suprâ umbrinâ, subtùs albicante, ferruginoso marginatâ tinctâque, fasciis nigricantibus duodenis retrices omnes transversim notantibus, binas autem intimas extimasque minùs conspicuè ; rhachibus desuper brunneis, subtùs albis : rostro cum unguibus nigro : cerâ, palpebris tibiisque olivascentibus : iride brunneâ.*

Head, back, wings, and tail, light blue ash-colour ; throat and under tail coverts white ; cervical collar, breast, belly, and thighs, pale rufous ; under parts of the wings, and the base of all the plumage, white. Wings rounded ; the fourth quill longest ; the first, half the length of the fourth ; shafts black ; tail slightly rounded, consisting of twelve feathers, the colour lightest beneath ; shafts black above, white beneath ; bill and claws blue-black ; cere, eyelids, and legs, rich gamboge-yellow ; irides orange.

The female does not differ from the male except in size, being much larger, and in having the under tail coverts pale rufous instead of white.

*Young.* Head and back umber, the feathers edged with ferruginous, and barred with white, the latter scarcely visible without raising the feathers ; throat white ; breast tawny-white, each of the feathers having a central line, and terminal spot of umber ; flanks barred with umber ; thighs tawny, with irregular bars of pale ferruginous ; wings umber, with darker bars ; beneath tawny, barred with umber ; shafts dusky beneath, brown above ; tail umber on the upper surface, whitish beneath, edged and tinged with ferruginous ; twelve dark bars cross the whole, but are palest on the two inner and two outer feathers ; shafts brown above, white beneath ; bill and claws black ; cere, eyelids, and legs, pale olive green ; irides brown. Young females agree with the above description of the male.