

3. Q U I N T I N I A, *Alph. DC.*1. Q U I N T I N I A S E R R A T A, *A. Cunn.*

*Quintinia serrata*, A. Cunn. Bot. N. Zeal. in Ann. Nat. Hist. 2, p. 256; Hook. Ic. Pl. t. 558; Hook. f. Fl. N. Zeal. p. 78.

HAB. New Zealand: in dry woods at the Bay of Islands. (In fruit.)

4. C A L L I C O M A, *Andr.*1. C A L L I C O M A S E R R A T I F O L I A, *Andr.*

*Callicoma serratifolia*, Andr. Bot. Rep. t. 566; Sims, Bot. Mag. t. 1811; DC. Prodr.

HAB. New South Wales; in the neighbourhood of Sydney.

5. C E R A T O P E T A L U M, *Smith.*1. C E R A T O P E T A L U M G U M M I F E R U M, *Smith.*

*Ceratopetalum gummiferum*, Smith, N. Holl. 1, p. 9, t. 3; DC. Prodr. 4, p. 13.

HAB. New South Wales, near Sydney.

6. S P I R Æ A N T H E M U M, *Nov. Gen.*

*Flores polygamo-dioici vel hermaphroditi. Calyx quadri-quinquefidus, æstivatione valvatus, persistens. Corolla nulla. Stamina 8 vel 10, imæ basi calycis inserta, fere hypogyna: filamenta filiformia, fl. masc. exserta, hermaphrodito-fert. calyce haud longiora: antheræ didymæ,*