

pezida. Prof. Williston's contribution to this volume, pp. 1-89, was published in 1891-1892, and the list of subsequently described species, and the general index to the whole of the three volumes, pp. 93-127, in 1903. The material gain was scanty in comparison with the large number of species which must exist in the region, but among the Syrphids the genera *Baccha*, *Volucella*, and *Eristalis* were particularly well represented. The total number of species enumerated is 325, of which 67 are new. The two plates include figures of 29 species.

46-47. **Rhynchota Heteroptera.** By W. L. Distant and G. C. Champion.
2 vols., pp. xx, 462, xvi, 416, with 16 plain, and 45 coloured plates.
£9 net

Vol. I. By W. L. Distant. *Pentatomidæ*, *Coreidæ*, *Lygæidæ*, *Pyrrhocoridæ*, and *Capsidæ*. Mr. Distant's enumeration of the above mentioned families of Rhynchota Heteroptera (pp. 1-303) was issued in 1880-1884, the Supplement to the same (pp. 304-351) in 1889-1893, and the Appendix (pp. 452-462) in 1893, the delay in publishing the Supplement being due to his long absence in South Africa. Altogether, 1108 species are recognized, more than half of which are described as new. In the Introduction, published in 1893, the number of genera belonging to each family is given and the geographical distribution shown in short tables. Amongst the Lygæidæ nearly half the genera are stated to be endemic. The Pentatomidæ include 377 species and are, therefore, very well represented in Central America, though many of the genera are common to the Neotropical region. The Capsidæ, too, with 313 species, are very numerous, but the tropical representatives of this family are but little known as yet, and no comparison with the fauna of the adjacent portion of South America can be made.

The thirty-nine plates include figures of upwards of 900 species, a complete list of them being given in the Introduction, pp. xi-xx.

Vol. II. By G. C. Champion. *Tingitidæ-Corixidæ*. Mr. Champion here deals with nineteen families of Rhynchota Heteroptera (making twenty-four in all), twelve belonging to the Gymnocerata and seven to the Cryptocerata. The number of species enumerated is 592, nearly half of which are described as new. Fifteen additions to vol. I., described or recorded by various authors in other works, between 1893 and 1901, are noted on p. 384, the total number of Heteroptera enumerated in the two volumes being 1715. The Tingitidæ, Aradidæ, Reduviidæ, and Anthocoridæ are abundantly represented in Central America, the respective numbers for each being 78 (66 new), 78 (40 new), 204 (78 new), and 54 (32 new). As with the five families belonging to vol. I., no comparison is made with the fauna of the adjacent regions southward for want of sufficient data. Most of the genera belonging to the aquatic