

7. *Cnemidophorus inornatus*.

Cnemidophorus inornatus, Baird, Proc. Ac. N. Sc. Philad. 1858, p. 255 ; Cope, Proc. Am. Phil. Soc. xvii. 1877, p. 93.

Hab. NORTH MEXICO, Nuevo Leon (*Baird*).

Four supraorbitals. Scales on the gular fold smaller than those on the breast anteriorly, and scarcely larger than those on the middle of the chin. Scales of the back tuberculated and elevated. Light greenish-olive, paler beneath. No lines on the body.

GYMNOPHTHALMUS.

Gymnophthalmus, Merrem, Tent. Syst. Amphib. p. 74.

Of the six species known, three are West-Indian, two South-, and one Central-American.

1. *Gymnophthalmus sumichrasti*.

Epaphelus sumichrasti, Cope, Journ. Ac. N. Sc. Philad. 1876, p. 115.

Gymnophthalmus sumichrasti, Bocourt, Miss. Sc. Mex., Rept. p. 471, t. 22 n. fig. 2.

Hab. MEXICO, Tehuantepec ; WEST GUATEMALA.—VENEZUELA (*Brit. Mus.*).

Sumichrast says that he has but rarely met with this lizard near Cacoprieto under rotting trunks in the forest.

TRETIOSCINCUS.

Tretioscincus, Cope, Proc. Ac. N. Sc. Philad. 1862, p. 184.

Beside the Nicaraguan species, only one other, from Colombia (*T. bifasciatus*, A. Dum.), is known.

1. *Tretioscincus lævicauda*.

Tretioscincus lævicauda, Cope, Proc. Am. Phil. Soc. xi. 1870, p. 557.

Hab. NICARAGUA.

ARGALIA.

Argalia, Gray, Ann. & Mag. Nat. Hist. 1846, xviii. p. 67.

1. *Argalia pæcilocheilus*.

Gerrhonotus pæcilocheilus, Lichtenst. Nomencl. Rept. Mus. Berol. p. 16.

Ecpleopus (Argalia) pæcilocheilus, Peters, Abhandl. Berl. Ak. 1863, p. 213.

Hab. PANAMA, Veragua.—VENEZUELA.