

### 9. *Gerrhonotus rudicollis*.

*Gerrhonotus rudicollis*, Wieg. Isis, 1828, p. 379; and Herpet. Mex. p. 31, t. 10. fig. 1; Bocourt, Miss. Sc. Mex., Rept. p. 367, t. 21 B. figg. 3.

*Barissia rudicollis*, Gray, Cat. Liz. p. 55.

*Hab.* MEXICO (*Deppe*).

Dorsal scales strongly keeled and, like the ventral, in twelve or fourteen, nuchal in four or six longitudinal series. Nasal in contact with the rostral; supranasals none. Three pairs of shields on the upper surface of the snout, between vertical and rostral. Sides with brown irregular cross bands.

One specimen only is known.

### 10. *Gerrhonotus imbricatus*.

*Gerrhonotus imbricatus*, Wieg. Isis, 1828, p. 379; and Herpet. Mex. p. 34, t. 10. figg. 2 & 5; Dum. & Bibr. Erp. Gén. v. p. 407; Bocourt, Miss. Sc. Mex., Rept. p. 363, t. 21 B. figg. 2.

*Gerrhonotus lichenigerus*, Wagl. Ic. Amph. t. 34. fig. 2; Wieg. Herpet. Mex. p. 35, t. 10. fig. 6 (*G. adspersus*); Bocourt, l. c. p. 365, t. 21 B. figg. 1 (*G. imbricata*, var.).

*Barissia imbricata*, Gray, Cat. Liz. p. 55.

*Hab.* MEXICO, Cuernavaca and Guanajuato (*Bocourt*), City of Mexico (*Brit. Mus.*), Vera Cruz (*Boucard*), Tehuantepec (*Sumichrast*).

Dorsal scales keeled, in twelve or fourteen longitudinal series. The nasal is not in contact with the rostral; supranasals none. Three pairs of shields on the upper surface of the snout, all very convex.

This well-known species is subject to variation of colour, white dots being irregularly scattered about the scutes of the head and body. The mental scute is generally followed by a pair of scutes, more rarely by a single one.

### 11. *Gerrhonotus planifrons*.

*Gerrhonotus planifrons*, Bocourt, l. c. p. 361, t. 21 c. figg. 1.

*Hab.* MEXICO, Oaxaca (*Boucard*).

This form, if it should prove to be a constant modification, differs from the typical *G. imbricatus* in having the head-shields flat, and sixteen longitudinal series of dorsal scales.

### 12. *Gerrhonotus cæruleus*. (Tab. XXV. figg. A-D.)

*Gerrhonotus cæruleus*, Wieg. Isis, 1828, p. 379; Dum. & Bibr. Erp. Gén. v. p. 397; Bocourt, Miss. Sc. Mex., Rept. p. 353, t. 21 c. figg. 3.

*Gerrhonotus burnettii*, Gray in Griff. An. King., Rept. p. 64; and in Beechey's Voy. Rept. p. 96, t. 31. fig. 2 (bad); and in Cat. Liz. p. 54; Bocourt, l. c. p. 356, t. 21 c. figg. 4.