

ENYALIUS.

Enyalius, Wagler, Syst. Amphib. p. 150.

Of the ten or eleven species of this South-American genus, only the following occur north of the Isthmus of Darien. We have not seen it.

1. *Enyalius heterolepis*.

Enyalius heterolepis, Bocourt, Ann. Sc. Nat. xix. 1874, art. 4, and Miss. Sc. Mex., Rept. t. 20 c. fig. 14.

Hab. PANAMA, Veragua.

IGUANA.

Iguana, Laurenti.

1. *Iguana rhinolophus*.

Iguana rhinolophus, Wiegman. Herp. Mex. p. 44; Dum. & Bibr. Erp. Gén. iv. p. 207; Tyler, Proc. Zool. Soc. 1850, t. 3 (*Metopoceros cornutus*); Cope, Proc. Ac. N. Sc. Phil. 1862, p. 356; Sumichrast, Ann. & Mag. N. H. 1864, xiii. p. 500; Bocourt, Miss. Sc. Mex., Rept. p. 134; Cope, Journ. Ac. N. Sc. Phil. viii. 1876, p. 124.

Hab. MEXICO, Mazatlan and Presidio (*Forrer*), Cordova (*Sallé*), Tehuantepec and Rio Coazacoalcos (*Sumichrast*), Cozumel Island on the coast of Yucatan (*G. F. Gaumer*); GUATEMALA, Chiapam (*Salvin*); COSTA RICA (*Cope*); PANAMA (*Brit. Mus.*), Darien, R. Truando (*Schott*).

"Iguana verde" of the Creoles; "Guchachi-guëla" of the Zapotec Indians.

This is the northern or Central-American race of the South-American *I. tuberculata*; it does not seem to extend southwards beyond the Isthmus of Darien, but occurs also in the island of Santa Lucia and probably other parts of the West Indies.

2. *Iguana tuberculata*.

Iguana tuberculata, Laur. Syn. Rept. p. 49.

Hab. GUATEMALA (*Salvin*); HONDURAS (*Brit. Mus.*); NICARAGUA (*Brit. Mus.*); PANAMA (*Dow*), Darien (*Schott*).—Generally distributed over the northern parts of SOUTH AMERICA and BRAZIL; occurring also in many West-Indian Islands.

CTENOSAURA.

Ctenosaura, Wiegmann, Isis, 1828, p. 371.

1. *Ctenosaura acanthura*.

Lacerta acanthura, Shaw, Zool. ii. 1802, p. 216.

Cyclura teres, Harlan, Journ. Ac. Sc. Phil. 1825, p. 246, t. 16.

Ctenosaura cycluroides, Wiegman. Isis, 1828, p. 371; Bocourt, Miss. Sc. Mex., Rept. p. 143 (part.); Sumichrast, Bull. Soc. Zool. Fr. v. 1880, p. 175.

Ctenosaura shawii, Gray, Wiegman.