

the neck smaller than those on the back. The large supraoculars not divided. Hind limb equal to the distance of the vent from the collar; fourth toe rather longer than the head. Femoral pores fifteen. Green or greenish-bronze coloured, without transverse collar on the neck; a large black patch on the collar-fold of the side of the neck, the two patches being connected by a black band across the throat. Young males have the abdomen blue, yellowish grey in the middle, the colours separated from each other by black. In very old males the light colour in the middle disappears, and the whole abdomen is uniform blackish blue. In an adult male each dorsal scale is black, with a broad green band, also the upperside of the head is symmetrically spotted with green.

Size of adult male 3 inches without tail.

Cope (*l. c.*) says that he has seen a specimen from Tehuantepec; but as this individual has a double row of supraoculars, the locality is not inserted here.

3. *Sceloporus poinsetti*.

Sceloporus poinsetti, Baird & Gir. Proc. Ac. N. Sc. Phil. vi. p. 126; and in U. S. & Mex. Bound. Surv., Rept. p. 5, t. 29. figg. 1-3.

Hab. NORTH AMERICA, Texas.—MEXICO, Rio San Pedro and Sonora (*Clark*), Ciudad in Durango (*Forrer*).

Twenty-one or twenty-two scales in a series along the back. The large supraoculars divided into a double series. Hind limb equal to the distance of the vent from the humeral joint; fourth toe as long as the head. Femoral pores eleven or twelve. Scales on the neck not conspicuously smaller than those of the back. A broad black collar edged with light green across the neck; body and tail with alternate dark and greenish cross-bands, the darker bands being the broader. Lower parts blue and black in the adult male, a whitish band along the middle of the abdomen.

Size of adult male 4 inches without tail.

The coloration of the upper parts is probably not constant.

4. *Sceloporus acanthinus*.

Sceloporus acanthinus, Bocourt, Miss. Sc. Mex., Rept. p. 180, t. 18. figg. 10, 10 *a*, *b*, & t. 19. figg. 4, 4 *a*; Boul. Cat. Liz. ii. p. 221.

Hab. MEXICO, Jalapa (*Höge*); GUATEMALA, San Agustin on the Volcan de Atitlan (*Mus. Paris*).

Twenty-four scales along the back. The large supraoculars not divided. Hind limb equal to the distance of the vent from the humeral joint; fourth toe longer than the head. From twelve to fourteen femoral pores. Anterior nuchal scales much smaller than those on the back. A narrow black collar only one or two scales broad across the neck.