

7. *Sceloporus undulatus*.

Agama undulata, Daud. Rept. iii. p. 384.

Sceloporus undulatus, Fitz. Syst. Rept. p. 75; Baird & Gir. U. S. Expl. Exped., Herp. p. 379, t. 19. figg. 15-21; Bocourt, Miss. Sc. Mex., Rept. p. 195, t. 18 bis. figg. 11, 11 *a*, *b*; Cope, Proc. Am. Phil. Soc. 1885, p. 398; Boul. Cat. Liz. ii. p. 227.

Sceloporus thayerii, Baird & Gir. Proc. Ac. N. Sc. Phil. vi. p. 127.

Hab. NORTH AMERICA, United States.—MEXICO, Sonora (*Baird & Girard*); GUATEMALA, Yzabal (*Salvin*).

From twenty-six to thirty-one scales in a series along the back. The large supraoculars not divided. Hind limb as long as, or shorter than, the distance of the vent from the collar; fourth toe as long or nearly as long as the distance of the snout from the lateral collar-fold. Thirteen to seventeen femoral pores. Greyish, brownish, or bronze-coloured, with irregular dark brown bands or transverse spots across the back; no nuchal collar.

An adult male is 3 inches long without tail.

This is a northern species which scarcely enters into the limits of the Central-American fauna. The reasons which induce me to admit it, are:—(1) because Baird and Girard say expressly in the description of their *Sceloporus thayerii* that it extends into Sonora; and (2) because Mr. Salvin's collection, made at Yzabal, contains a specimen which, although it has only twenty-six scales in the dorsal series, cannot be kept distinct from *S. undulatus*.

8. *Sceloporus lunæi*.

Sceloporus lunæi, Bocourt, Miss. Sc. Mex., Rept. p. 184, t. 18 bis. figg. 5, 5 *a*, *b*.

Sceloporus formosus, part., Boul. Cat. Liz. ii. p. 222.

Hab. GUATEMALA, Coban (*Salvin*), plateau 1500 metres (*Bocourt*).

Twenty-five scales in a series along the back, the dorsal series distinctly convergent on the posterior part of the back; scales keeled, with a long terminal spine. A single series of large supraoculars. One canthal scale. The anterior lobes of the ear-opening not larger than the preceding scales. Hind limb nearly as long as the distance of the vent from the collar; fourth toe longer than the head. Femoral pores thirteen. Green; each dorsal scale with a yellowish spot at the base; blue underneath, the middle of the abdomen being yellowish with a black streak on each side; collar-fold deep black, this colour not extending across the neck or throat.

Size of adult male 3 inches without tail.

9. *Sceloporus irazuensis*, sp. n.

Sceloporus formosus, part., Boul. Cat. Liz. ii. p. 222.

Hab. MEXICO, Southern States?—COSTA RICA, Irazu (*Rogers*).