

with yellow; abdomen whitish. Male brownish, faintly marbled with darker colour, sides marbled with yellowish; a narrow yellow vertebral line and a light interorbital cross-band. Female nearly uniform brownish-black above; sides marbled with yellowish.

	♀.	♂.
Length of the body	60 millim.	51 millim.
Distance between vent and heel	62 „	52 „
Length of hind limb (from vent)	104 „	87 „

I have long hesitated before I ventured to distinguish this frog under a distinct name; however, the receipt of a second specimen agreeing in every essential distinctive character with the first, removed every doubt about the species being distinct from *L. fleischmanni*. The male has the testicles well developed, yet lacks vocal sacs. These organs are present in much smaller specimens of *L. fleischmanni*.

6. *Liohyla engytympanum*, sp. n. (Tab. LXV. fig. A.)

Hab. COSTA RICA, Monte Redondo (*Underwood, in Mus. Brit.*).

In general habit and form of the head similar to *L. rugulosa*. Snout rather obtuse, with the canthus rostralis distinct, and with sloping sides; eye scarcely equal to its distance from the nostril. *Tympanum* an irregular oval, deeper than long, barely half the area of the orbit (in adult female). Vomerine teeth in two short prominent transverse series, slightly separated from each other, behind but within the level of the choanæ; choanæ rather wider than the eustachian openings. Upper and lateral parts of the body, supra-orbital region, and upper parts of the hind limbs with tubercles and granules; a glandular fold above the tympanum; abdomen smooth, with indistinct granules on the sides; transverse and longitudinal folds indicate the presence of an abdominal disk. Length of the body a little less than the distance between vent and heel. Disks of fingers and toes subequal, rounded, moderately developed; first finger rather longer than the second; a rudimentary web between the three inner fingers. Metatarsals without pads, except at the base of the inner toe; outer metatarsal tubercle scarcely visible; a tarsal fold. Toes one-fourth webbed. Greyish, with a yellow vertebral and gular line; an indistinct dark interorbital cross-band. Upper lip and flanks with blackish spots. Upper parts of the limbs obscurely banded; hind part of the femur rather finely and conspicuously marbled with black and white. Lower parts white.

One adult female.

Length of the body	59 millim.
Distance between vent and heel	62 „
Length of foot	47 „