

2. *Hyla bocourti*.

Hyla regilla, Brocchi, Miss. Sc. Mex., Batr. p. 31, t. 13. fig. 2 (not B. & G.).

Hyliola bocourti, Mocquard, Nouv. Arch. Mus. i. p. 341 (1899); Bull. Soc. Philom. (9) i. p. 164 (1899).

Hab. MEXICO, Tepic (*Diguet*); GUATEMALA, Alta Vera Paz (*Bocourt*).

Closely allied to *H. eximia*, but with the posterior face of the thigh ornamentally coloured.

Vomerine teeth between the choanæ, in two transverse groups; choanæ and eustachian ostia nearly equal. Snout of moderate length, rather depressed, with obtuse canthus and sloping sides. Distance of the nostril from the eye nearly as long as the diameter of the latter, and more than the distance of the nostrils from each other. Interorbital space wider than the upper eyelid. Tympanum distinct, one-half the area of the eye. Fingers free, with disks somewhat larger than those of the toes, and somewhat smaller than the tympanum. Toes barely one-third webbed; (tarsal fold, if present, quite indistinct). Upper parts smooth, lower granular. Upper parts bluish, with large blackish longitudinal spots, which commence between the eyes. Side of the snout blackish, which colour is continued as a band across the tympanum and along the side of the body; this band is sharply defined towards the back, and finely edged with white, but towards the abdomen it gradually passes in an irregular outline into the light coloration of the lower parts; it has also an incomplete whitish lower edge between the eye and the tympanum. Upper jaw without a horn-coloured border; limbs indistinctly cross-banded, and the posterior face of the thigh is finely marbled with black and white on a greyish ground.

			♀.
Greatest width of head			15 millim.
Distance of vent from snout			37 „
„ „ heel			33 „
„ „ metatarsal joint			43 „
„ „ end of fourth toe			59 „

This description is taken from one of the specimens collected by Bocourt, and kindly communicated to me by Professor Vaillant; I have deposited it in the Natural History Museum.

3. *Hyla nana*, sp. n. (Tab. LXXIII. fig. A.)

Hab. MEXICO, Cuernavaca (*H. H. Smith*).

Vomerine teeth (sometimes indistinct or even entirely absent) between the choanæ, in two fascicles; choanæ wider open than eustachian ostia. Snout of moderate length, rather pointed, not depressed, with angular canthus and subvertical sides. Eye of moderate size, its diameter equal to its distance from the nostril, and more than the