

Light grey, with five darker longitudinal bands, the median of which expands anteriorly so as to form a large subtriangular spot between the eyes; femur with one, tibia with three cross-bands; concealed surfaces unspotted. (After Cope.)

Length of body	29 millim.
„ hind limb	43 „
„ hind foot	19 „

13. *Hyla phæota*.

Hyla phæota, Cope, Proc. Ac. N. Sc. Philad. 1862, p. 358; Werner, SB. Bayer. Akad. Wiss. xxvii. p. 97 (1897).

Hyla labialis, Peters, MB. Ak. Wiss. Berlin, 1863, p. 463; 1873, p. 618.

Hab. PANAMA (*Werner*), Chiriqui (*Peters*).—COLOMBIA, Turbo (*Cope*), Bogota (*Peters*).

Vomerine teeth in two transverse series between the large choanæ. Head broad, depressed, with rounded canthus rostralis. Tympanum more than half the extent of the eye. The three outer fingers slightly webbed; nearly three phalanges of the fourth toe free, the web, however, margining the outer side. The tibio-tarsal articulation reaches well beyond the end of the snout, when the hind limb is carried forwards. Upper parts smooth; a strong fold from the eye over the tympanum. Brownish-olive above, with a longitudinal band or series of spots on the back, bifurcating anteriorly; a spot on the coccyx and near the vent brown. Upper jaw, postorbital region, and the margins of the upper eyelid and of the limbs white. A dark band along the canthus rostralis is continued over the tympanum to the scapular region; femur with narrow cross-bands and with the hinder side reticulate or uniformly coloured; tibiæ more broadly cross-banded; sides with brown vermiculations; lower parts whitish. Male with an unpaired vocal sac. (After Cope.)

Length	22 lines.
------------------	-----------

The descriptions of other authors who believe that they have recognized Cope's species differ in several noteworthy particulars. Thus, Peters states that "the tympanum is scarcely visible, its diameter being two-fifths of the diameter of the eye"; the web between the toes is deeply excised, but "otherwise nearly complete, only the last phalanx of the fourth toe being free." On the other hand, Werner describes the tympanum as "equal to $\frac{3}{4}$ of the diameter of the eye. Fingers with $\frac{1}{3}$, toes with nearly complete webs"; he also ascribes to this species a tarsal fold.

14. *Hyla tæniopus*, sp. n. (Tab. LXXII. fig. F.)

Hab. MEXICO, Jalapa (*Trujillo*).

Vomerine teeth in two transverse series, between the choanæ. Head broad, depressed, broader than the body, with the eyes obliquely directed forwards; snout short, not