

24. *Hyla underwoodi*.

*Hyla microcephala*, Bouleng. P. Z. S. 1898, p. 481, t. 39. fig. 3 (nec Cope).

*Hyla underwoodi*, Bouleng. Ann. & Mag. N. H. 1899, iii. p. 277.

*Hab.* COSTA RICA, Bebedero (*Underwood*).

Vomerine teeth in two small groups well separated from each other, between the choanæ. Head rather small, a little longer than broad; snout not depressed, with angular canthi, which convergent form a somewhat acute angle in front; the distance between the nostrils is much less than their distance from the orbit, or than the diameter of the eye. Tympanum distinct, one-fourth the area of the eye. Fingers one-third, toes three-fourths webbed; disks of both subequal, and about the size of the tympanum. *No tarsal fold*. Upper parts smooth; abdomen granular; no folds. Pinkish, powdered with brown; a brown band, edged with white above, runs along the side of the snout, and through the eye to the side of the body, gradually fading behind; the white edge marks distinctly the canthus and margin of upper eyelid, and is continued above the tympanum across the shoulder. Thighs not pigmented. Lower parts white.

Greatest width of head . . . . .	8.5 millim.
Distance of vent from snout . . . . .	28.5 „
„ „ heel . . . . .	28 „
„ „ metatarsal joint . . . . .	34 „
„ „ end of fourth toe . . . . .	46 „

25. *Hyla nigripes*.

*Hyla nigripes*, Cope, Journ. Ac. N. Sc. Philad. viii. p. 104, t. 23. fig. 7 (1876).

*Hab.* COSTA RICA (*Cope*).

Vomerine teeth between the choanæ, in two short transverse series. *Shape of head oval, tongue longer than broad*; canthus rostralis a little concave; sides of the snout oblique. Tympanum one-fourth or one-fifth the area of the eye; eustachian ostia and choanæ nearly equal. Fingers with the web extending to about one-half the length of the longest; web of toes not extending to the end of the penultimate phalanx; disks of moderate size. Smooth above, finely areolate below. Dark brown, limbs and feet blackish; groin finely white and blackish marbled; no lateral border or band. Posterior face of femur black; no spots on any other region. (After Cope.)

Width of head . . . . .	13 millim.
Length of body . . . . .	39 „
„ hind limb . . . . .	65 „
„ hind foot . . . . .	16 „