

26. *Hyla molitor*.

Hyla molitor, Schmidt, Denkschr. Ak. Wiss. Wien, xiv. p. 425, t. 1. figg. 8, 9 (1858); Peters, MB. Ak. Wiss. Berlin, 1873, p. 618.

Hab. PANAMA, Chiriqui Lagoon, near Boca del Toro.

Vomerine teeth between the choanæ, in two groups well separated from each other. Snout with distinct strongly convergent canthi and with sloping sides. Tympanum small, obliquely elliptical; eustachian ostia nearly as wide as the choanæ. The three outer fingers with a rudimentary web; toes fully webbed to the disks. Upper parts smooth; no fold across the chest; an indistinct one above the tympanum. Upper parts greyish blue, uniform or marbled with black; a series of black dots along the canthus rostralis; sides and thighs generally marbled with blackish. (After Schmidt.)

Length of body	33 millim.	38 millim.
„ hind limb	53* „	60 „

27. *Hyla miotympanum*.

Hyla miotympanum, Cope, Proc. Ac. N. Sc. Philad. 1863, p. 47; Bull. U. S. Nat. Mus. no. 32, p. 14 (1887); Peters, MB. Ak. Wiss. Berlin, 1871, p. 651.

Hyla microtis, Peters, l. c. 1869, p. 880.

Hab. MEXICO, Jalapa and Zacualtipan (*Cope*), Puebla (*Berkenbusch*), Mirador (*Sartorius*), Vera Cruz and Tehuantepec (*Sumichrast*).

Vomerine teeth in two slightly oblique series converging backwards between the choanæ, which are much wider than the eustachian ostia. Snout broadly rounded, without distinct canthus, and with sloping sides; nostrils as far distant from each other as from the eyes. *Tympanum extremely small, scarcely visible*. Fingers one-half webbed or less, with disks which are larger than the tympanum. Toes broadly webbed. Upper parts smooth, lower granular; no folds or appendages. Purplish above, with or without a few lighter specks and indistinct darker reticulations; a yellow band along the margin of the upper jaw, continued on the side of the body, this portion sometimes not well separated from the abdominal shade. Limbs not barred; concealed portions of the thigh not coloured. (After Cope and Peters.)

Width of head	13 millim.
Length of body	37 „
„ hind limb	60 „

Cope describes the lateral band as yellow. Peters states: "On the boundary between the dark coloration of the back and the lighter of the abdomen there is a narrow line or a punctated series of black colour"; he does not mention the lateral band.

* 33 in Schmidt's description is obviously a misprint.