

28. *Hyla prosoblepon*.

*Hyla prosoblepon*, Böttger, Katal. Batr. Senckenb. Mus. p. 45 (1892).

*Hylella puncticrus*, Bouleng. Ann. & Mag. N. H. 1896, xviii. p. 341.

*Hab.* COSTA RICA, Limon (*Fleischmann*), La Palma (*Underwood*); PANAMA, Bugaba (*Champion*).

Vomerine teeth (if present) in two transverse series, between the choanæ. Head broad, depressed, with very short snout, and with *large eyes, directed forwards*; canthus obtuse, concave, sides of the snout sloping. Diameter of the eye a little more than its distance from the nostril, and much more than the distance between the nostrils; upper eyelid narrow, narrower than the interorbital space. Tympanum very small, deeply sunk, about one-sixth the area of the eye. The two outer fingers one-third webbed, but the third and second are connected only by a rudiment of a web; toes four-fifths webbed. The disks of the fingers are larger than those of the toes and than the tympanum. When carried forwards, the heel reaches beyond the end of the snout. Upper parts smooth, abdomen coarsely areolated. Light purplish above with scattered round dark purple or blackish dots; lower parts whitish; the concealed parts of the limbs not pigmented, the remainder coloured like the back. Humerus of the male with a projecting, curved, compressed osseous process.

			♀.
Width of head . . . . .			9 millim.
Distance of vent from snout . . . . .			25 „
„ „ heel . . . . .			27 „
„ „ metatarsal joint. . . . .			35 „
„ „ end of fourth toe . . . . .			42 „

A small species, in which the vomerine teeth appear to be as frequently absent as present; specimens with and without them occur in the same locality.

29. *Hyla pulverata*.

*Hyla pulverata*, Peters, MB. Ak. Wiss. Berlin, 1873, p. 614.

*Hab.* PANAMA, Chiriqui.

Vomerine teeth in two groups between the choanæ, which are much wider than the narrow eustachian ostia. Snout as long as the eye, with obtuse canthus. *Tympanum very small and covered with pigmented skin*. The three outer fingers broadly webbed, the web extending to the disks of the second and fourth fingers, but not to that of the third; toes broadly webbed. Upper parts smooth, abdomen granulated; no fold above the tympanum. Colourless, with scattered white points. (After Peters.)

Length of body . . . . .	24 millim.
Width of head . . . . .	8 „
Length of hind limb . . . . .	45 „
„ foot . . . . .	20 „