

Hab. PANAMA, Chiriqui (*Schmidt*).—TROPICAL SOUTH AMERICA.

Vomerine teeth between, and extending to behind, the wide choanæ, in two oblique, arched series: (\ . Snout broad and depressed, with rather indistinct canthus and sloping sides. Distance of the nostril from the eye equal to the diameter of the latter, but more than the distance between the nostrils. Interorbital space wider than the supraocular cover. Tympanum three-fourths the area of the eye. The three outer fingers one-third, more rarely one-fourth webbed, their disks equal to those of the toes, but much smaller than the tympanum. A conspicuously projecting rudiment of the pollex. Toes three-fourths webbed, with moderate subarticular tubercles; large specimens with a slight tarsal fold. Upper parts smooth; a transverse fold or series of tubercles above the vent. Abdomen with the proximal part of the throat granular. Upper parts marbled with brown or uniform whitish or pinkish; sides with narrow vertical blackish bars; hinder part of the thigh with similar bars, which extend across towards the front, though much fainter. Male with a subgular vocal sac.

	♀.	♂.
Greatest width of head	24 millim.	20 millim.
Distance of vent from snout	70 „	60 „
„ „ heel	75 „	64 „
„ „ metatarsal joint	93 „	83 „
„ „ end of fourth toe	115 „	105 „

37. *Hyla albomarginata*.

Hyla albomarginata, Spix, Spec. Nov. p. 33, t. 8. fig. 1; Dum. et Bibr. Erpét. Gén. viii. p. 555; Günth. Batr. Sal. p. 98; Peters, MB. Ak. Wiss. Berlin, 1872, p. 207; Bouleng. Batr. Sal. p. 356.

Hypsiboas albomarginata, Wagl. Syst. Amphib. p. 201; Cope, Proc. Am. Phil. Soc. 1886, p. 273.

Hyla infulata, Wied, Beitr. Naturg. Bras. i. p. 533; Burmeister, Erläut. Faun. Bras. p. 97, t. 30. figg. 1-6.

Hyla punctata, Wied, Abbild. tab. —.

Hab. NICARAGUA (*Bransford*); COSTA RICA, San Carlos (*Underwood*).—TROPICAL AMERICA.

Vomerine teeth between the wide choanæ, in two rather long oblique series, generally arched: (\ . Snout broad and depressed, with somewhat pointed nose, rather indistinct canthus, and sloping sides. Distance of the nostril from the eye a little more than the diameter of the latter. Interorbital space much wider than the upper eyelid. Tympanum distinct, one-third the area of the eye. The three outer fingers are more than half-webbed, and provided with disks at least as large as the tympanum, and larger than the disks of the toes. Toes almost entirely webbed, with moderate subarticular tubercles; no tarsal fold. Upper parts smooth; generally a very distinct fold, which starts from the eye and descends towards the middle of the