

middle of dorsal fin. Dorsal IX 10 ; second spine longest,  $\frac{2}{3}$  the length of head. Anal III 7 ; second spine not longer than third, about  $\frac{2}{5}$  the length of head. Pectoral longer than the head. Caudal deeply forked. Silvery, back brownish ; about 10 narrow bluish vertical bars on the upper part of the side : dorsal and caudal fins dusky.

*Hab.* Pacific Coast from Lower California to Panama :  
Rio Presidio, Mexico.

This species differs slightly from its Atlantic representative (*G. cinereus*) in the fewer scales, larger head, longer maxillary, and longer second anal spine, whilst the bars on the sides are usually more numerous.

3. **Gerres gula**, Cuv. & Val., 1830.

*Eucinostomus gula*, Jord. & Everm. t. c. p. 1370.

*Hab.* Atlantic Coast from Carolina to Brazil.

The top of the head is covered with scales to the level of the nostrils except for a small rounded or oval naked area completely surrounded by scales, overlying the præmaxillary spines.

4. **Gerres harengulus**, Goode & Bean, 1879.

*Eucinostomus harengulus*, Jord. & Everm. t. c. p. 1368.

*Hab.* Atlantic Coast from Florida to Bahia.

5. **Gerres californiensis**, Gill, 1862.

*Eucinostomus dowi*, Jord. & Everm. t. c. p. 1367.

*Eucinostomus californiensis*, Jord. & Everm. t. c. p. 1369.

*Hab.* Pacific Coast from California to Panama :  
Rio Mascota, Mexico (*Buller*).

6. **Gerres pseudogula**, Poey, 1875.

*Eucinostomus pseudogula*, Jord. & Everm. t. c. p. 1368.

*Hab.* Atlantic Coast from the Bermudas to Brazil.

7. **Gerres lefroyi**, Goode, 1874.

*Ulæma lefroyi*, Jord. & Everm. t. c. p. 1371.

*Eucinostomus meeki*, Eigenm. Bull. U.S. Fish. Comm. xxii. 1902, p. 229, fig. 10 (1904).

*Hab.* Atlantic Coast from the Bermudas to Cuba ; rivers of Cuba.

8. **Gerres peruvianus**, Cuv. & Val., 1830.

*Gerres peruvianus*, Jord. & Everm. t. c. p. 1376.

*Hab.* Pacific Coast from Mexico to Peru ; Rio Guayas, Ecuador.