

Hab. HONDURAS (*Dyson* ^{1 2 3 4 5}).

Remarkable for its slender turreted form and neat coloration variegated with white and dark reddish-brown.

Doubtful Species of Cyclostomidæ.

Chondropoma truncatum.

Cyclostoma truncatum (Wiegman), Rossm. Icon. vi. p. 49, t. 28. fig. 397 (1839) ¹; v. Mart. in Malak. Blätt. xiii. p. 5 (1865) ².

Chondropoma truncatum, Pfr. Monogr. Pneum. Vivent. i. p. 293 ³; Gray, Cat. Phaneropn. p. 208 ⁴.

Hab. MEXICO ^{1 2 3 4} (*Deppe*).

The original specimen is lost. The author's description and figure agree rather well with *Chondropoma turritum*, Pfr.; but this has hitherto only been known from Honduras, and Deppe did not collect in that country, but in Mexico.

Tudora planospira.

Cyclostoma (Tudora) planospira, Pfr. in Malak. Blätt. iii. p. 208 (1856) ¹.

Tudora planospira, Pfr. Monogr. Pneum. Vivent. ii. p. 127 ².

Hab. "In republica Mexicana, teste Poey."

No other species of *Tudora* is known from the American continent. No figure is given.

DIPLOMMATINA.

Diplommatica, Benson, Ann. & Mag. Nat. Hist. iv. (2), p. 193 (1849).

Shell ovate, thin, with vertical riblets; whorls somewhat irregular, the one before the last the largest; aperture vertical, circular; peristome double, expanded. Operculum thin, shelly, paucispiral. Tentacles long, filiform; eyes bilobated, sessile at the hinder part of the base of the tentacles.

1. Diplommatica stollii, sp. n. (Tab. I. figg. 19 a, b.)

"Eine winzige Art, welche einen sonst blos aus Ostindien bekannten linksgewundenen starkgerippten Typus darstellt."—*O. Stoll*, Guatemala, Reisen und Schilderungen aus den Jahren 1878–1883, p. 199 (1886).

Testa sinistrorsa, imperforata, ovato-conica, costulis in anfr. penultimo 16 sat validis, albida; anfr. 6, convexi, priores duo læves, flavescentes, penultimus maximus, ultimus paullo angustior, rotundatus, læviusculus; apertura subverticalis, circularis, peristomate continuo, incrassato, expanso, albo.

Long. $2\frac{1}{2}$, diam. 1, apert. $\frac{2}{3}$ millim.