

1. CENTRAL MEXICO, N. of the tropic.

Helicina durangoana and *H. borealis*.

2. CENTRAL MEXICO, between the tropic and the Isthmus of Tehuantepec.

Helicina ghiesbreghti and *H. tenuis*.

3. WESTERN SLOPE OF MEXICO, between the same limits.

Helicina punctisulcata, *H. fragilis*.

4. EASTERN SLOPE OF MEXICO, between the same limits, chiefly State of Vera Cruz.

Helicina cinctella, *H. zephyrina* (*turbinata*), *H. deppeana*, *H. cordilleræ*, *H. chrysocheila*, *H. tenuis*, *H. fragilis*, *H. succincta*, *H. delicatula*, *H. oweniana*, var., *H. notata*, *H. flavida*, *H. lirata*, *H. mohriana*.

Schasicheila nicoleti, *S. pannucea*, *S. alata*.

Proserpina eolina, *P. salleana*, *P. berendti*.

This large number is due chiefly to the circumstance that many collectors, and among them some very zealous and exact, have collected in this district for a considerable time.

5. SOUTH-EASTERN MEXICO, Chiapas, Tabasco.

Helicina ghiesbreghti, *H. tenuis*, *H. oweniana*, *H. florida*, *H. lirata*.

6. YUCATAN.

Helicina tenuis, *H. arenicola*.

7. BRITISH HONDURAS.

Helicina dysoni.

8. NORTH-WEST GUATEMALA, Valley of the R. Usumacinta below Salinas.

Helicina amœna, *H. tenuis*, *H. fragilis*, *H. oweniana*, var., *H. flavida*.

Schasicheila pannucea.

9. NORTH-EAST GUATEMALA, Valley of the R. Polochic and of the R. Usumacinta above Salinas.

Helicina amœna, ? *H. sowerbyana*, *H. rostrata*, *H. tenuis*, *H. fragilis*, *H. oweniana*, var., *H. flavida*, *H. chryseis*, *H. microdina*.

Schasicheila pannucea.

10. SOUTH GUATEMALA, Pacific slope.

Helicina ghiesbreghti, *H. tenuis*, *H. lirata*.

Schasicheila pannucea.