

16. ***Helicina succincta***, sp. n. (Tab. I. figg. 6; 9, var.)

Testa globoso-conoidea, leviter spiratim sulcata, albida vel pallide rufescens, fusco-variegata; anfr. 5, convexiusculi, sutura impressa, ultimus prope aperturam basi planulatus, angulo valde obtuso prope aperturam evanescente cinctus; apertura diagonalis, triangularem-rotundata, peristomate incrassato, breviter expanso, albo, margine columellari subperpendiculari, in tuberculum subrectangulum terminato, callo crasso, parvo, circumscripto.

Diam. maj. 9-10, min. $7\frac{1}{2}$ -9, alt. $8\frac{1}{2}$ -10 millim.; apert. diam. 4- $4\frac{1}{2}$, alt. obliqua 5-6 millim. (fig. 6).

Hab. E. MEXICO: Cordova (*Sallé, Höge*) (fig. 6).

Var. minor, diam. 8, alt. $6\frac{1}{2}$ millim., rufescens, fusco-strigata (fig. 9).

Hab. E. MEXICO: Cuesta de Misantla (*Mateo Trujillo*) (fig. 9); Tlacolula, between Las Vigas and Misantla (*Höge*).

Rather near *H. arenicola*, especially as regards the form of the peristome and columella; but of larger size, with more elevated spire, remarkably coarser spiral structure, and more reddish coloration. As the localities are also well separated, and I have seen no intermediate forms, I feel obliged to keep them distinct; it is possible, however, that when additional examples are obtained they may prove to run one into the other.

17. ***Helicina raresulcata***.

Helicina raresulcata, Pfr. in Malak. Blätt. viii. p. 173 (1861)¹; Monogr. Pneum. Vivent. iii. p. 228², & iv. p. 269³; Strebel, Abhandl. Geb. Naturw. Ver. Hamb. vi. 1, p. 19, t. 1 a. fig. 9, & t. 2. figg. 9, 9 a⁴.

Hab. E. MEXICO: Vera Cruz (*Berendt*^{1 2}); Vera Cruz, on the plain inland of the sandhills, on the leaves and stalks of a stinging shrub called "mala muger" (*Strebel*⁴).

18. ***Helicina arenicola***.

Helicina arenicola, Morelet, Test. Noviss. i. p. 21¹; Pfr. in Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, *Helicina*, no. 99, p. 73, t. 10. figg. 24-27²; Monogr. Pneum. Vivent. i. p. 366³; Gray, Cat. Phaneropn. p. 265⁴; Sow. Thes. Conch. iii. p. 286, t. 271. fig. 303⁵; Reeve, Conch. Icon. xix., *Helicina*, t. 28. fig. 250⁶.

Hab. N. YUCATAN: Sisal, in sandy localities^{3 4} (*Morelet*¹).

The habitat "Cuba" quoted by Sowerby⁵ and Reeve⁶ is erroneous; Morelet described in the same pamphlet¹ several Cuban species, but to this he expressly attributes the locality given above.

19. ***Helicina punctisulcata***, sp. n. (Tab. I. fig. 10.)

Testa conoidea, subangulata, sulcis tenuibus subdistantibus punctiferis sculpta, nitidula, rufescens, interdum fusco-fasciata, subtus pallidior; spira conica, rufa; apertura diagonalis, triangularem-semicircularis, peristomate incrassato, leviter expanso, albo, margine columellari brevi, subperpendiculari, basi subangulato, callo parvo, flavo.

Diam. maj. 9, min. $7\frac{1}{2}$, alt. $7\frac{1}{2}$ millim.; apert. diam. $4\frac{1}{2}$, alt. obliqua 4 millim.