

22. *Helicina oweniana*. (Tab. I. figg. 11, 12.)

a. genuina: subelate conoidea, pallida, peristomate aurantio; subinde unifasciata (fig. 12).

Helicina oweniana, Pfr. in P. Z. S. 1848, p. 123¹; Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, *Helicina*, no. 48, p. 40, t. 7. figg. 35, 36²; Monogr. Pneum. Vivent. i. p. 373³ (excl. var.) & ii. p. 199; Gray, Cat. Phaneropn. p. 270⁴; Sow. Thes. Conch. iii. p. 287, t. 272. figg. 241, 242⁵; Reeve, Conch. Icon. xix., *Helicina*, t. 16. figg. 144⁶.

Hab. S.E. MEXICO: Chiapas (*Ghiesbreght*^{1 2 3 4}); Teapa in Tabasco (*H. H. Smith*).

b. coccinostoma: globoso-conoidea, alba, citrina vel fulva absque fasciis, peristomate coccineo.

Helicina coccinostoma, Morelet, Test. Noviss. i. p. 19 (1849)⁷; Sow. Thes. Conch. iii. p. 289, t. 273. figg. 295, 296⁸; Reeve, Conch. Icon. xix., *Helicina*, t. 20. fig. 173 (magnified)⁹.

Helicina oweniana, var. β , Pfr. Monogr. Pneum. Vivent. i. p. 373¹⁰; Gray, Cat. Phaneropn. p. 270¹¹.

Hab. N. GUATEMALA: Province of Peten¹¹ (*Morelet*^{7 10}).

c. anozona: subglobosa, rubescens vel fulvescens, zona suturali pallida, peristomate luteo (fig. 11).

Helicina oweniana, Tristr. in P. Z. S. 1861, p. 233¹².

Helicina anozona, v. Mart. in P. Z. S. 1875, p. 649¹³.

Hab. N. AND CENTRAL GUATEMALA: Coban (*Salvin*^{12 13}); Cubilguitz, Northern Vera Paz (*Champion*); Telemán, in the valley of the Polochic River (*Stoll*).

These three forms can be well distinguished in the majority of specimens; but there are also some others which are intermediate between them, either in colour or in shape. All those I have seen have a similar fine sculpture.

23. *Helicina notata*.

Helicina notata (Sallé), Pfr. in P. Z. S. 1856, p. 323, t. 35. figg. 18-20¹; Monogr. Pneum. Vivent. ii. p. 203²; Sow. Thes. Conch. iii. p. 287, t. 272. figg. 239, 240³; Reeve, Conch. Icon. xix., *Helicina*, t. 31. fig. 276⁴.

Hab. E. MEXICO: Cordova, State of Vera Cruz (*Sallé*^{1 2}); Atoyac (*Höge*); Mexico^{3 4}.

Easily recognizable by a peculiar dark red spot on the back of the penultimate whorl above the suture, the rest of the shell being unicolorous yellow. The specific name probably refers to this mark; but it is not even mentioned in Pfeiffer's description, although visible in the specimens in his collection. The malleated surface is somewhat like that of many specimens of *H. tenuis*; the spiral ridges bring it near to *H. succincta*.

24. *Helicina flavida*.

Helicina flavida, Menke, Synops. Moll. ed. 1, p. 79 (1828)¹, ed. 2, p. 132 (1830)²; Sow. Thes. Conch. i. p. 9, t. 3. figg. 117, 134³, & iii. p. 287, t. 272. figg. 233-236⁴; Pfr. in Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, *Helicina*, no. 52, p. 42, t. 1. figg. 31, 32, & t. 5. figg. 28-