

VI. *Depressæ*.26. *Helicina borealis*, sp. n. (Tab. I. fig. 15.)

Testa depressa, sublævis, striatula, alba, superne maculis griseis biseriatis plus minusve obsoletis picta; spira breviter conoidea, apice prominulo; anfr. 5, ultimus rotundatus, infra paullo magis convexus; apertura parva, valde semicircularis, peristomate incrassato, brevissime expanso, margine columellari brevi, subperpendiculari, in angulum prominulum terminata, callo crasso, circumscripto.

Diam. maj. 10-12, min. $8\frac{1}{2}$ -9, alt. 7 millim.; apertura diam. $4\frac{1}{2}$ -5, lat. $5-5\frac{1}{2}$ millim.

Hab. N. CENTRAL MEXICO: Villa Lerdo in Durango, in an open country with patches of *Mimosæ* (Höge).

Approaches in form *H. guadelupensis*, Sow.

27. *Helicina sanguinea*.

Helicina sanguinea, Pfr. in P. Z. S. 1848, p. 124¹; Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, *Helicina*, no. 10, p. 17, t. 8. figg. 20, 21²; Pfr. Monogr. Pneum. Vivent. i. p. 345³; Gray, Cat. Phaneropn. p. 249⁴; Sow. Thes. Conch. iii. p. 292, t. 275. figg. 364, 365⁵; Reeve, Conch. Icon. xix., *Helicina*, t. 20. figg. 178⁶.

Hab. HONDURAS^{5 6} (*Dyson*^{3 4}).

28. *Helicina dysoni*.

Helicina dysoni, Pfr. in P. Z. S. 1848, p. 121¹; Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, *Helicina*, no. 63, p. 49, t. 3. figg. 36-39²; Pfr. Monogr. Pneum. Vivent. i. p. 384³; Gray, Cat. Phaneropn. p. 278⁴.

Hab. VENEZUELA: Cumana (*Dyson*^{1 3}).

Var. a: minor.

Helicina dysoni, vars. β - δ , Pfr. Monogr. Pneum. Vivent. i. p. 384⁵; Martini & Chemnitz, Syst. Conch.-Cab. ed. 2, no. 63, p. 49, t. 8. figg. 27-32⁶; Sow. Thes. Conch. iii. p. 286, t. 271. figg. 213, 214⁷.

Hab. HONDURAS⁷ (*Dyson*⁵).

Var. b: subglobosa.

Helicina bocourti, Crosse & Fischer, in Journ. de Conch. xvii. p. 251 (1869)^{*}

Hab. BRITISH HONDURAS: Belize (*Bocourt*⁸).

Var. c, jansoni, v. Mart.: distincte angulosa (diam. maj. 5, alt. 4, apert. 2 millim.) (Tab. I. fig. 16.)

Hab. HONDURAS: Island of Bonacca, Bay of Honduras (*Gaumer*).

The type is stated to be from Cumana, Venezuela, by Dr. Pfeiffer^{1 3}. According to Mr. Bland the same species occurs also on the island of Trinidad, on the coast of Venezuela, and has been described by J. L. Guppy [Ann. & Mag. Nat. Hist. (3) xiv.