

6. *Glandina aurata*.

Glandina aurata, Morelet, Test. Noviss. i. p. 12 (1849)¹; Fischer & Crosse, Miss. Scient. Mex., Mollusca, i. p. 106, t. 3. figg. 7, 7 a²; Angas, P. Z. S. 1879, p. 480, t. 40. fig. 6 (anim.)³; Jousseau, Bull. Soc. Zool. Fr. iii. p. 169 (1878)⁴.

Achatina (*Glandina*) *aurata* (Morelet), Pfr. Monogr. Helic. Vivent. iii. p. 517⁵.

Oleacina aurata (Morelet), Gray, Cat. Pulm. p. 33⁶; Pfr. Monogr. Helic. Vivent. iv. p. 642⁷.

Achatina lignaria, Reeve, Conch. Icon. vi., *Achatina*, t. 8. fig. 27⁸; Pfr. Monogr. Helic. Vivent. iii. p. 519⁹.

Oleacina lignaria (Reeve), Gray, Cat. Pulm. p. 34¹⁰; Pfr. Monogr. Helic. Vivent. iv. p. 642¹¹.

Glandina lignaria (Reeve), H. & A. Adams, Gen. Moll. ii. p. 108¹²; v. Mart. P. Z. S. 1875, p. 647¹³.

Glandina sowerbyana, Form C, Strebel, Beitr. Mex. Land- und Süßw.-Conch. ii. p. 17, t. 6 a. figg. 12, 12 a-f¹⁴.

Glandina, sp., Binney, Ann. N. Y. Acad. i. p. 261 (1879)¹⁵; iii. p. 81, t. 17. fig. E (1884) (radula)¹⁶.

Hab. N. GUATEMALA: Woods of Vera Paz (*Morelet*¹); Alta Vera Paz (*Bocourt*²); Vera Paz^{5 6} (*Boucard*⁴); Coban (*Salvin*¹³).

CENTRAL GUATEMALA: Mountain forest between Tepan and Totonicapam 8000 to 9000 feet above the sea (*Stoll*).

S.E. COSTA RICA (*Gabb*^{3 15 16}).

Strebel's *G. aurata* (op. cit. ii. p. 21, t. 12. figg. 36, 36 a, b), from Ocaña, Colombia, belongs, I believe, to another species, distinct by the more turrite-cylindrical form, the shorter aperture, the want of the numerous dark brown stripes, and a somewhat weaker sculpture; a specimen collected by Herr Alf. Stübel in the Llanos of San Martin appears to be referable to the same species, for which I propose the name *G. stübeli*.

The locality for Reeve's specimen of *A. lignaria* was unknown⁸.

B. *Marginatae*.

7. *Glandina fusiformis*.

Achatina fusiformis, Pfr. P. Z. S. 1845, p. 75¹; Monogr. Helic. Vivent. ii. p. 292²; Reeve, Conch. Icon. vi., *Achatina*, t. 9. fig. 31³; Deshayes, in Férussac's Hist. Nat. Moll. Terr. ii. p. 181, t. 137. figg. 1-3⁴.

Glandina fusiformis, Morelet, Journ. de Conch. iii. p. 33 (1852)⁵; Fischer & Crosse, Miss. Scient. Mex., Mollusca, i. p. 103, t. 3. fig. 2 a⁶; Strebel, Beitr. Mex. Land- und Süßw.-Conch. ii. p. 26, t. 9. figg. 14, 14 a⁷; v. Mart. P. Z. S. 1875, p. 647⁸; Jousseau, Bull. Soc. Zool. Fr. iii. p. 169 (1878)⁹.

Oleacina fusiformis (Pfr.), Gray, Cat. Pulm. p. 32¹⁰; Pfr. Monogr. Helic. Vivent. iv. p. 642¹¹.

Var. *miltochila*: anfr. ultimis stramineis, peristomate miniaceo.

Achatina fusiformis, Reeve, loc. cit. fig. 31 b.

Glandina fusiformis, var. β , Fischer & Crosse, loc. cit. p. 104, t. 3. fig. 2.

BIOL. CENTR.-AMER., Terr. and Fluvial. Mollusca, April 1891.